arXiv:1003.2004v1 [math.PR] 10 Mar 2010

Submitted to the Annals of Applied Probability

ON THE RATE OF CONVERGENCE TO STATIONARITY OF THE M/M/N QUEUE IN THE HALFIN-WHITT REGIME

By David Gamarnik and David Goldberg

MIT

We study the rate of convergence to stationarity of the $M/M/n$ queue in the Halfin-Whitt regime. We prove that there is an interesting *phase transition* in the system's behavior, occurring when the excess parameter B reaches $B^* \approx 1.85772$. For $B < B^*$, the exponential rate of convergence is $\frac{B^2}{4}$; above B^* it is the solution to an equation involving the parabolic cylinder functions. We also bound the prefactor governing the rate of convergence uniformly over n when $B < B^*$, and use our bounds to derive a rule-of-thumb for determining the time it takes a severely overloaded (underloaded) queueing system to return (probabilistically) to the steady-state.

1. Introduction. Parallel server queueing systems can operate in a variety of regimes that balance between efficiency and quality of offered service. This is captured by the Halfin-Whitt (H-W) regime, which can be described as critical with respect to the probability that an arriving customer has to wait for service. Namely, in this regime the stationary probability of wait is bounded away from both 0 and 1 as the number of servers grows. Although studied originally by Erlang [16] and Jagerman [25], the regime was formally introduced in [22] by Halfin and Whitt, who established the criticality of the probability that a customer waits in terms of the square-root spare capacity rule for large multiserver queues with exponentially distributed service times. They also showed that as the size of the multiserver queue grows, the queueing process (properly scaled) converges weakly to a non-trivial positive recurrent diffusion (the H-W diffusion). Similar weak-convergence results were subsequently obtained for more general multiserver systems [34], [26], [32], [19], with the most general results appearing in [36]. This line of research has been motivated by questions related to the management of call centers, since it is generally accepted that well-run call centers operate in the H-W regime [1],[20].

Since the steady-state behavior of the $M/M/n$ queue in the H-W regime is quite simple in practice [22], while the transient dynamics are more complicated [22], it is common to use the steady-state approximation to the transient distribution. Thus it is important to understand the *quality* of the steady-state approximation. The only work along these lines seems to be the recent paper [30], which studies the transform of the H-W diffusion and proves several results analogous to our own for the H-W diffusion. The key difference is that in this paper we study the pre-limit $M/M/n$ queue, not the limiting diffusion. We note that the relevant transform functions were also studied in [2], although in a different context.

The question of how quickly the positive recurrent $M/M/n$ queue approaches stationarity has a rich history in the queueing literature. In [33], Moorse derives an explicit solution for the transient $M/M/1$ queue, and discusses implications for the exponential rate of convergence to stationarity. Similar analyses are carried out in [10] and [37]. Around the same time, two different research

---

AMS 2000 subject classifications: Primary 60K25

Keywords and phrases: Many-server queues, rate of convergence, spectral gap, weak convergence, orthogonal polynomials

groups [31], [27] worked out powerful and elegant theories that could be used to give the transient distributions for large classes of birth-death processes (b-d-p). The transient probabilities are expressed as integrals against a *spectral measure* $\phi(x)$ that is intimately related to the eigenvalues of the generator of the b-d-p. Karlin and McGregor (K-M) devote an entire paper [28] to the application of their theory to the $M/M/n$ queue. For a positive recurrent b-d-p $Z(t)$, let $P_{i,j}(t) \overset{\Delta}{=} \Pr(Z(t) = j | Z(0) = i)$, $P(t) \overset{\Delta}{=} (P_{i,j}; i, j \in Z^+)$, and $P_j(\infty) \overset{\Delta}{=} \lim_{t\to\infty} P_{i,j}(t)$ denote the steady-state distribution. Then K-M proved

THEOREM 1. *[28] For any positive recurrent $M/M/n$ or $M/M/\infty$ queue, there exist a set of polynomials $\{Q_i(x)\}$, a set of strictly positive constants $\{c_i\}$, and a measure $\phi(x)$ with non-negative support such that*

$$P_{i,j}(t) = c_j \int_0^\infty e^{-xt} Q_i(x) Q_j(x) d\phi(x). \tag{1}$$

K-M also comment on the relationship between the rate of convergence to stationarity and the support of $\phi(x)$ [28]. This relationship was later made rigorous and expanded on in a series of papers [4],[5],[40],[6]. Recall that the spectral gap γ of a b-d-p is the absolute value of the supremum of the set of strictly negative real eigenvalues of the associated generator matrix A, where $\frac{d}{dt}P(t) = AP(t)$ [17], [6]. If no such eigenvalues exist, we set $\gamma = \infty$. For a complex-valued function $f(x)$, we let $Z(f)\big(Z^+(f)\big)$ denote the infimum of the set of (strictly positive) real zeros of $f(x)$. Set $Z(f)\big(Z^+(f)\big) = \infty$ if $f(x)$ has no (strictly positive) real zeros. We then have

THEOREM 2. *[28],[4],[5],[40],[6] For any positive recurrent $M/M/n$ or $M/M/\infty$ queue, $\gamma \in (0, \infty)$. For all i and j, $\lim_{t\to\infty} -t^{-1} \log |P_{i,j}(t) - P_j(\infty)| \geq \gamma$. For at least one pair of i, j, $\lim_{t\to\infty} -t^{-1} \log |P_{i,j}(t) - P_j(\infty)| = \gamma$. Furthermore, $\gamma = Z^+(\phi)$.*

It is well-known that for the positive recurrent $M/M/1$ and $M/M/\infty$ queues, γ can be computed explicitly. In particular,

THEOREM 3. *[28] For the positive recurrent $M/M/1$ queue with arrival rate λ and service rate μ, $\gamma = (\sqrt{\lambda} - \sqrt{\mu})^2$, and the* spectral measure *$\phi(x)$ consists of a jump at 0, and an absolutely continuous measure on $[(\sqrt{\lambda} - \sqrt{\mu})^2, (\sqrt{\lambda} + \sqrt{\mu})^2]$. For the $M/M/\infty$ queue with arrival rate λ and service rate μ, $\gamma = \mu$, and the* spectral measure *$\phi(x)$ consists of a countably infinite number of jumps, at the points $\{k\mu; k \in Z^+\}$.*

Unfortunately, for the general positive recurrent $M/M/n$ queue, the known characterizations for γ are cumbersome and hard to use. The difficulty is that for the positive recurrent $M/M/n$ queue with arrival rate λ and service rate μ, the *spectral measure* $\phi(x)$ consists of three parts [28]. The first part is a jump at 0, which corresponds to the steady-state distribution [28]. The second component is an absolutely continuous measure on the interval $[(\sqrt{\lambda} - \sqrt{n\mu})^2, (\sqrt{\lambda} + \sqrt{n\mu})^2]$, whose density is described in [28]. The third component consists of a set of at most n (but possibly zero) jumps, which all exist on $\big(0, (\sqrt{\lambda} - \sqrt{n\mu})^2\big)$ [28]. The complexity of determining γ arises from the difficulty of locating these jumps [40]. In [28], the set of jumps is expressed in terms of the zeros of a complicated polynomial equation.

Significant progress towards understanding these jumps was made in a series of papers by van Doorn [38],[39],[40],[41]. Van Doorn used the K-M representation and the theory of orthogonal polynomials to give several alternate characterizations and bounds for the spectral gap of a b-d-p, and applied these to the $M/M/n$ queue. He also showed that for each fixed n there is a transition in the nature of the spectral measure of the $M/M/n$ queue as one varies the traffic intensity [38].

THEOREM 4. *[38] For all $n \geq 1$, there exists $\rho_n^* \in [0,1)$ such that for any $M/M/n$ queue satisfying $\frac{\lambda}{n\mu} \geq \rho_n^*$, $\gamma = (\sqrt{\lambda} - \sqrt{n\mu})^2$; and for any $M/M/n$ queue satisfying $\frac{\lambda}{n\mu} < \rho_n^*$, $\gamma < (\sqrt{\lambda} - \sqrt{n\mu})^2$.*

Unfortunately, all of the characterizations (including that of ρ_n^*) given by van Doorn are again fairly complicated, and van Doorn himself comments in [40] that one is generally better off using the approximations that he gives in the same paper. Van Doorn's work was later extended in [29], and similar results were achieved using different techniques in [46]. It was also shown in [46] that $\rho_n^* \leq (1 - \frac{1}{n})^2$.

There are also some results in the literature for explicitly bounding the distance to stationarity (as opposed to just identifying the exponential rate of convergence) [46],[7]. These bounds are generally given in terms of an explicit prefactor attached to an exponentially decaying term (often containing the spectral gap). However, these bounds are generally not studied in the context of weak convergence, and thus may not scale desirably with n. In Section 8 we will show that two such bounds appearing in the literature ([46] and [7]) do not scale properly in the H-W regime. We note that the complexity of bounding the distance to stationarity uniformly for a sequence of b-d-p is related to the cutoff phenomenon for Markov chains [11], which has been studied in the context of queueing systems [18]. We also note that the asymptotic properties of the bounds appearing in [46] have recently been analyzed for several models different from those studied in this paper [42],[43].

1.1. *Outline of paper.*

In Section 2, we introduce some notation and state our main results. In Section 3, we prove some technical preliminaries, cite several results from [29] and [38], and prove a new characterization for γ_n (Lemma 3). Sections 4 - 7 are devoted to studying the asymptotic properties of Lemma 3, with each section analyzing a different term appearing in Lemma 3. This culminates with Section 7, in which the asymptotic results of Sections 4 - 6 are combined with the characterization for γ_n given in Lemma 3 to prove our main results (Theorem 5, Corollary 1, and Theorem 6). In Section 8 we prove our explicit bounds on the distance to stationarity (Theorem 7), and compare to other bounds from the literature. In Section 9 we summarize the main results of the paper and present ideas for future research. We note that several of our proofs necessitate a considerable amount of simple algebra, and we have generally chosen to include the details, with an eye towards making the proofs easier to understand in their totality.

2. Main Results.

2.1. *Definitions and notations.*

We now define several important quantities for the $M/M/n$ queue with arrival rate $\lambda_n \triangleq n - B\sqrt{n}$ and service rate $\mu \triangleq 1$. We assume that $B > 0$, and n is large enough to ensure that $\lambda_n > 0$ and $n > \lambda_n + 1$. Let $Q(t)$ denote the queue length at time t, namely the total number of customers including those in service. Let $P^n_{i,j}(t) \triangleq \Pr(Q(t) = j | Q(0) = i)$, $P^n_j(\infty) \triangleq \lim_{t\to\infty} P^n_{i,j}(t)$, $P^n_{i,\leq j}(t) \triangleq \sum_{k=0}^{j} P^n_{i,k}(t)$, and $P^n_{\leq j}(\infty) \triangleq \sum_{k=0}^{j} P^n_k(\infty)$. Let γ_n denote the spectral gap of the associated Markov chain. We let $H_k \triangleq \sum_{j=1}^{k} \frac{1}{j}$ denote the kth harmonic number, and $\zeta \triangleq \lim_{n\to\infty} \big(H_n - \log(n)\big) \approx .5772$ denote the Euler-Mascheroni constant. All logarithms will be base e. Unless otherwise stated, all functions are defined only for real values of x. All empty products are assumed to be equal to unity, and all empty summations are assumed to be equal to zero.

2.2. *The parabolic cylinder functions.* We now briefly review the two-parameter function commonly referred to as the parabolic cylinder function $D_x(z)$, since we will need these functions for the statement (and proofs) of our results. For excellent references on these functions, see [21] Section 8.31 and Section 9.24, [3] Sections 3.3-3.5, and [15] Chapter 8. Let $\Gamma(x)$ denote the Gamma function (see [23], Chapter 8.8). It is stated in [3] that for real x and z, $D_x(z) \in \mathbb{R}$, and

$$D_x(z) = \begin{cases} \sqrt{\frac{2}{\pi}} e^{\frac{z^2}{4}} \int_0^\infty e^{-\frac{t^2}{2}} Cos(\frac{\pi}{2}x - zt) t^x dt & \text{if } \ x \geq 0; \\ \frac{e^{-\frac{z^2}{4}}}{\Gamma(-x)} \int_0^\infty e^{-\frac{t^2}{2} - zt} t^{-(x+1)} dt & \text{if } \ x < 0. \end{cases} \tag{2}$$

$D_x(z)$ takes on a simpler form for $x \in Z$. In particular, it is stated in [21] that for $z \in \mathbb{R}$,

$$D_{-1}(z) = \sqrt{2} e^{\frac{z^2}{4}} \int_{\frac{z}{\sqrt{2}}}^{\infty} e^{-t^2} dt, \quad D_0(z) = e^{-\frac{z^2}{4}}, \quad \text{and} \quad D_1(z) = z e^{-\frac{z^2}{4}}. \tag{3}$$

Note that since $\Gamma(-x) \in (0, \infty)$ for $x < 0$, (2) and (3) imply that

$$\text{for all} \quad z \in \mathbb{R} \quad \text{and} \quad x \leq 0, \quad D_x(z) > 0. \tag{4}$$

The parabolic cylinder functions arise in several contexts associated with the limits of queueing models, such as the Ornstein-Uhlenbeck limit of the appropriately scaled infinite server queue [24] and various limits associated with the Erlang loss model [44]. We note that the parabolic cylinder functions have been studied as the limits of certain polynomials under the H-W scaling, using tools from the theory of differential equations [12],[13],[14],[35].

2.3. *Main results.* We now state our main results. We begin by describing the asymptotic phase transition that occurs w.r.t. the spectral gap γ_n of the $M/M/n$ queue in the H-W regime. Let

$$\zeta(B) \triangleq \begin{cases} \frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} + \frac{B}{2} & \text{if } D_{\frac{B^2}{4}-1}(-B) \neq 0, \\ \infty & \text{otherwise;} \end{cases} \tag{5}$$

and

$$\Psi_\infty(x) \triangleq \begin{cases} \frac{D_x(-B)}{D_{x-1}(-B)} + \frac{1}{2}\big(B + (B^2 - 4x)^{\frac{1}{2}}\big) & \text{if } D_{x-1}(-B) \neq 0, x \leq \frac{B^2}{4}; \\ \infty & \text{otherwise.} \end{cases} \tag{6}$$

Note that $\zeta(B) = \Psi_\infty(\frac{B^2}{4})$. We include a plot of $\zeta(B)$.



Fig 1. $\zeta(B)$

Let $B^* \stackrel{\Delta}{=} Z^+(\zeta)$. Then

Proposition 1. *$B^* \in [\sqrt{2}, 2)$, and $Z^+(\Psi_\infty) < \min(1, \frac{B^2}{4})$ for $B > B^*$. Numerically, $B^* \approx 1.85772$.*

Our main result is that

Theorem 5. *The limit $\lim_{n\to\infty} \gamma_n \stackrel{\Delta}{=} \gamma_B$ exists for all $B > 0$. For $0 < B \le B^*$, $\gamma_B = \frac{B^2}{4}$. For $B \ge B^*$, $\gamma_B = Z^+(\Psi_\infty)$.*

We include a plot of γ_B.



Fig 2. γ_B

We note that due to the highly non-linear manner in which the steady-state probability of wait scales in the H-W regime (see [22]), the case $0 < B < B^*$ actually encompasses most scenarios of interest. Indeed, it is proven in [22] that $\lim_{n\to\infty} P^n_{\leq n}(\infty) = 1 - \big(1 + B\exp(\frac{1}{2}B^2)\int_{-\infty}^{B}\exp(-\frac{1}{2}z^2)dz\big)^{-1}$. As this limit is monotone in B, the case $0 < B < B^*$ includes all scenarios for which the steady-state probability of wait lies in the set $\Bigg(\big(1 + B^*\exp(\frac{1}{2}(B^*)^2)\int_{-\infty}^{B^*}\exp(-\frac{1}{2}z^2)dz\big)^{-1}, 1\Bigg) \approx (.038, 1)$.

The following corollary may be interpreted as an asymptotic version of Theorem 4, unifying several earlier characterizations for the spectral gap of the $M/M/n$ queue [28],[38],[40],[46],[29].

Corollary 1. *The ρ_n^* parameter of Theorem 4 satisfies*

$$\lim_{n\to\infty}\sqrt{n}(1-\rho_n^*) = B^*.$$

The following theorem lists some properties of γ_B.

Theorem 6. *γ_B is a continuous, strictly increasing function of B on $(0,\infty)$. $\lim_{B\to\infty}\gamma_B = 1$, and $\lim_{B\to\infty}\frac{\log(1-\gamma_B)}{B^2} = -\frac{1}{2}$.*

We now give an interpretation of Theorem 5 and Corollary 1. The $M/M/n$ queue behaves like an $M/M/1$ queue when all severs are busy, and an $M/M/\infty$ queue when at least one server is idle. The phase transition of Theorem 5 formalizes this relationship in a new way. For $0 < B < B^*$, the K-M spectral measure of the $M/M/n$ queue in the H-W regime has no jumps away from the origin, and has spectral gap equal to $(\sqrt{\lambda_n} - \sqrt{n})^2$, two properties shared by the associated $M/M/1$ queue (see Theorem 3). For $B > B^*$, the K-M spectral measure has at least one jump away from the origin, like the associated $M/M/\infty$ queue (whose spectral measure has only jumps and spectral gap equal to 1, see Theorem 3). Another interpretation is that the $M/M/n$ queue cannot approach stationarity faster than either component system would on its own.

We now give our explicit bounds on the distance to stationarity for the case $B < B^*$.

Theorem 7. *For all $B \in (0, B^*)$ and $a_1, a_2 \in \mathbb{R}$, let $\alpha = \max(B, |a_1|, |a_2|)$. Then there exists N_{B,a_1,a_2} s.t. $n \geq N_{B,a_1,a_2}$ implies that for all $t \geq 1$,*

$$|\sqrt{n}P^n_{\lceil n+a_1\sqrt{n}\rceil,\lceil n+a_2\sqrt{n}\rceil}(t) - \sqrt{n}P^n_{\lceil n+a_2\sqrt{n}\rceil}(\infty)| \leq 2(\pi t)^{-\frac{1}{2}}(1+n^{-\frac{1}{8}})\exp\big(2(\alpha+B)^2 + 18\alpha^2 t^{-1} - \frac{B^2}{4}t\big),$$

and

$$|P^n_{\lceil n+a_1\sqrt{n}\rceil,\leq\lceil n+a_2\sqrt{n}\rceil}(t) - P^n_{\leq\lceil n+a_2\sqrt{n}\rceil}(\infty)| \leq 4(B^2\pi t)^{-\frac{1}{2}}(1+n^{-\frac{1}{8}})\exp\big(2(\alpha+B)^2) + 18\alpha^2 t^{-1} - \frac{B^2}{4}t\big).$$

Note that Theorem 7 provides a bound for any sufficiently large *fixed* n and *all times* greater than 1, which is stronger than the bounds that would follow by naively applying the weak-convergence theory. Indeed, using the weak-convergence theory one could derive bounds of the form 'for any *fixed* time t, there exists a sufficiently large N_t s.t. $n \geq N_t$ implies ...', but these bounds would be too weak to make statements about *all time* for any *fixed* n. We note that although we were able to derive partial results for the case $B \geq B^*$, the derived bounds were considerably more complicated than those of Theorem 7, and we leave it as an open question to derive simple explicit bounds for the case $B \geq B^*$.

We now discuss a practical 'rule-of-thumb' interpretation of Theorem 7 in the context of managing call centers. Suppose one is operating a call center in the H-W regime (with $0 < B < B^*$) and at time $t = 0$ the center begins either very overloaded ($Q(0) = n + a\sqrt{n}$) or underloaded ($Q(0) = n - a\sqrt{n}$) with $a \gg 0$ a large constant (independent of n). Then Theorem 7 says that for any $\epsilon > 0$, the probability that a caller at time $t^*_{a,B} \stackrel{\Delta}{=} \frac{8}{B^2}a^2 + \frac{4}{B^2}log(\frac{1}{\epsilon})$ has to wait for service will be roughly within ϵ of the probability that a caller in steady-state has to wait for service. In particular, after $t^*_{a,B}$ time units have elapsed, the system will have returned to within ϵ of steady-state.

3. Technical Preliminaries .

In this section we introduce several notations, functions, and results from the literature that will be critical for our analysis. We then give a new characterization for γ_n (Lemma 3), which will be amenable to asymptotic analysis. We also prove some basic asymptotic results that will be used throughout the paper, and state some additional properties of the parabolic cylinder functions.

We begin by associating several functions to the $M/M/n$ queue, as in [29] and [38]. Let

$$f_{n,k}(x) \stackrel{\Delta}{=} \begin{cases} 1 & \text{if k = 0;} \\ 1 + \lambda_n - x & \text{if k = 1;} \\ (\lambda_n + k - x) f_{n,k-1}(x) - \lambda_n (k-1) f_{n,k-2}(x) & \text{otherwise.} \end{cases}$$

By a simple induction argument, for $0 \le k \le n$,

$$f_{n,k}(x) = \sum_{j=0}^{k} \binom{k}{j} \lambda_n^j \prod_{i=1}^{k-j} (i - x). \tag{7}$$

Also note that

$$Z(f_{n,n-1}) > 1, \quad \text{and} \quad f_{n,n-1}(x) > 0 \quad \text{for} \quad x < Z(f_{n,n-1}). \tag{8}$$

Indeed, for $x \le 1$ and $0 \le j \le n-1$, $\prod_{i=1}^{k-j}(i - x) \ge 0$, and by (7), $f_{n,n-1}(x)$ is a non-negative sum of such terms with the $(n-1)$st term (λ_n^{n-1}) strictly positive. Let

$$a_n(x) \stackrel{\Delta}{=} \begin{cases} \frac{1}{2}\Big(\lambda_n + n - x - \big((\lambda_n + n - x)^2 - 4\lambda_n n\big)^{\frac{1}{2}}\Big) & \text{if } x \le (\sqrt{n} - \sqrt{\lambda_n})^2; \\ \infty & \text{otherwise.} \end{cases}$$

Note that $a_n(x)$ is real-valued for $x \le (\sqrt{n} - \sqrt{\lambda_n})^2$, and therefore

$$a_n(x) = \frac{1}{2}\Big(\lambda_n + n - x - \big((\sqrt{n} - \sqrt{\lambda_n})^2 - x\big)^{\frac{1}{2}} \big((\sqrt{n} + \sqrt{\lambda_n})^2 - x\big)^{\frac{1}{2}}\Big). \tag{9}$$

Also,

$$a_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) = \sqrt{\lambda_n n}. \tag{10}$$

We also define

$$a_\infty(x) \stackrel{\Delta}{=} \begin{cases} \frac{1}{2}\big(B - (B^2 - 4x)^{\frac{1}{2}}\big) & \text{if } x \le \frac{B^2}{4}; \\ \infty & \text{otherwise.} \end{cases}$$

Let

$$b_n(x) \triangleq \begin{cases} \left(4\lambda_n n - (\lambda_n + n - x)^2\right)^{\frac{1}{2}} & \text{if } (\sqrt{n} - \sqrt{\lambda_n})^2 \leq x \leq (\sqrt{n} + \sqrt{\lambda_n})^2; \\ \infty & \text{otherwise.} \end{cases}$$

Note that $b_n(x)$ is real-valued for $x \in [(\sqrt{n} - \sqrt{\lambda_n})^2, (\sqrt{n} + \sqrt{\lambda_n})^2]$, and therefore

$$b_n(x) = \left(x - (\sqrt{n} - \sqrt{\lambda_n})^2\right)^{\frac{1}{2}} \left((\sqrt{n} + \sqrt{\lambda_n})^2 - x\right)^{\frac{1}{2}}. \tag{11}$$

Let $\sigma_n(y) \triangleq f_{n,n}(y) - \sqrt{\lambda_n n} f_{n,n-1}(y)$, $\psi_n(y) \triangleq f_{n,n}(y) - a_n(y) f_{n,n-1}(y)$,

$$z_{n,k}(y) \triangleq \begin{cases} \frac{f_{n,k}(y)}{f_{n,k-1}(y)} & \text{if } f_{n,k-1}(y) \neq 0; \\ \infty & \text{otherwise;} \end{cases}$$

and $z_n(y) \triangleq z_{n,n}(y)$. We also define

$$z_\infty(y) \triangleq \begin{cases} \frac{D_y(-B)}{D_{y-1}(-B)} + B & \text{if } D_{y-1}(-B) \neq 0; \\ \infty & \text{otherwise;} \end{cases}$$

Let

$$\Psi_n(y) \triangleq \begin{cases} z_n(y) - a_n(y) & \text{if } z_n(y) \neq \infty \text{ or } a_n(y) \neq \infty; \\ \infty & \text{otherwise;} \end{cases}$$

and

$$Q_{n,k}(x) \triangleq \begin{cases} 1 & \text{if k} = 0; \\ 1 - \frac{x}{\lambda_n} & \text{if k} = 1; \\ (1 - \frac{x}{\lambda_n} + \frac{\min(k-1,n)}{\lambda_n}) Q_{n,k-1}(x) - \frac{\min(k-1,n)}{\lambda_n} Q_{n,k-2}(x) & \text{otherwise.} \end{cases} \tag{12}$$

We note that the $Q_{n,k}(x)$ polynomials can be associated with the well-studied Poisson-Charlier polynomials (see the discussion in [38]). It is shown in [38] that

$$x \in \left((\sqrt{n} - \sqrt{\lambda_n})^2, (\sqrt{n} + \sqrt{\lambda_n})^2\right) \text{ implies } c_n(x) \triangleq Q^2_{n,n}(x) - Q_{n,n-1}(x) Q_{n,n+1}(x) > 0. \tag{13}$$

Let

$$g_n(k) \triangleq \begin{cases} \frac{\lambda_n^k}{k!} & \text{if } \ 0 \leq k \leq n; \\ \frac{\lambda_n^n}{n!} (\frac{\lambda_n}{n})^{k-n} & \text{otherwise.} \end{cases}$$

We now cite several results from [29] and [38] that characterize γ_n and $P^n_{i,j}(t)$.

Theorem 8. *[29] If $Z(\sigma_n) \geq (\sqrt{n} - \sqrt{\lambda_n})^2$, then $\gamma_n = (\sqrt{n} - \sqrt{\lambda_n})^2$. If $Z(\sigma_n) < (\sqrt{n} - \sqrt{\lambda_n})^2$, then $\gamma_n = Z(\psi_n)$.*

Theorem 9. *[28], [38] If $\frac{\lambda_n}{n} \geq \rho^*_n$, then for all i, j,*

$$P^n_{i,j}(t) - P^n_j(\infty) = (2\pi)^{-1} g_n(j) \frac{n!}{\lambda_n^n} (\lambda_n n)^{-1} \int_{(\sqrt{n} - \sqrt{\lambda_n})^2}^{(\sqrt{n} + \sqrt{\lambda_n})^2} e^{-xt} Q_{n,i}(x) Q_{n,j}(x) b_n(x) c_n(x)^{-1} dx. \tag{14}$$

We now cite some properties of $z_n(x), a_n(x)$, and $\Psi_n(x)$ stated in [29], all of which follow from elementary calculus and simple induction arguments, or the interlacing property of the zeros of orthogonal polynomials [8].

LEMMA 1. *[29]*

- *(i)* *$Z(f_{n,k-1}) \geq Z(f_{n,k})$ for $k \leq n$.*
- *(ii)* *For $x \in \big(-\infty, Z(f_{n,n-1})\big)$ and $k \leq n$, $z_{n,k}(x)$ is a strictly positive, continuous, and strictly decreasing function of x.*
- *(iii)* *For $x \in \big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$, $a_n(x)$ is a strictly positive, continuous, strictly increasing function of x.*
- *(iv)* *For $x \in \Big(-\infty, \min\big((\sqrt{n}-\sqrt{\lambda_n})^2, Z(f_{n,n-1})\big)\Big)$, $\Psi_n(x)$ is a continuous, strictly decreasing function of x. Also, if $(\sqrt{n}-\sqrt{\lambda_n})^2 < Z(f_{n,n-1})$, then $\Psi_n(x)$ is continuous at $(\sqrt{n}-\sqrt{\lambda_n})^2$.*

We now prove some additional properties of $\{z_{n,k}(x)\}$.

LEMMA 2. *$z_{n,k}(x)$ is a twice-differentiable concave function of x on $(0,1)$ for all $k \leq n$.*

PROOF. We proceed by induction on k. For the base case, consider $k = 1$. By definitions, $z_{n,1}(x) = \lambda_n + 1 - x$, and the assertion is trivial. Now, let us assume the statement is true for $j = 1, \ldots, k-1$ with $k-1 \leq n-1$. By definitions, $f_{n,k}(x) = (\lambda_n + k - x) f_{n,k-1}(x) - \lambda_n (k-1) f_{n,k-2}(x)$. Since by (8), $Z(f_{n,n-1}) > 1$, we have by Lemma 1.i that $f_{n,k-1}(x) > 0$ and $f_{n,k-2}(x) > 0$, from which it follows that $z_{n,k-1}(x) > 0$ and $z_{n,k}(x) = (\lambda_n + k - x) - \lambda_n (k-1)(z_{n,k-1}(x))^{-1}$. Thus

$$
\begin{aligned}
\frac{d^2x}{dx^2} z_{n,k}(x) &= -\lambda_n (k-1) \frac{d^2x}{dx^2}\big(z_{n,k-1}(x)^{-1}\big) \\
&= -\lambda_n (k-1) \frac{d}{dx}\Big(-z_{n,k-1}(x)^{-2} \frac{d}{dx} z_{n,k-1}(x)\Big) \\
&= \lambda_n (k-1)\Big(-2 z_{n,k-1}(x)^{-3}\Big(\frac{d}{dx} z_{n,k-1}(x)\Big)^2 + z_{n,k-1}(x)^{-2} \frac{d^2x}{dx^2} z_{n,k-1}(x)\Big). \qquad (15)
\end{aligned}
$$

Since $z_{n,k-1}(x) > 0$, $\big(\frac{d}{dx} z_{n,k-1}(x)\big)^2 > 0$, and by the induction hypothesis $\frac{d^2x}{dx^2} z_{n,k-1}(x) \leq 0$, it follows from (15) that $z_{n,k}(x)$ is twice-differentiable on $(0,1)$ and satisfies $\frac{d^2x}{dx^2} z_{n,k}(x) \leq 0$ (concavity), proving the induction. ☐

We now prove a slight variant of Theorem 8, which will be more amenable to asymptotic analysis.

LEMMA 3.

- *(i)* *If $(\sqrt{n}-\sqrt{\lambda_n})^2 < 1$ and $\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) < 0$, then $Z(\Psi_n) = Z^+(\Psi_n)$ is the unique zero of $\Psi_n(x)$ in the interval $\big(0, (\sqrt{n}-\sqrt{\lambda_n})^2\big)$ and $\gamma_n = Z^+(\Psi_n)$.*
- *(ii)* *If $(\sqrt{n}-\sqrt{\lambda_n})^2 < 1$ and $\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) \geq 0$, then $\gamma_n = (\sqrt{n}-\sqrt{\lambda_n})^2$.*
- *(iii)* *If $(\sqrt{n}-\sqrt{\lambda_n})^2 \geq 1$, then $Z(\Psi_n) = Z^+(\Psi_n)$ is the unique zero of $\Psi_n(x)$ in the interval $\big(0,1\big)$ and $\gamma_n = Z^+(\Psi_n)$.*

PROOF. We begin by proving some properties of Ψ_n and σ_n. From (7),

$$\begin{aligned}
\Psi_n(0) &= \frac{\sum_{k=0}^{n}\binom{n}{k}\lambda_n^k\prod_{i=1}^{n-k} i}{\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{i=1}^{n-1-k} i} - \frac{1}{2}\Big(\lambda_n + n - \big((\lambda_n+n)^2 - 4\lambda_n n\big)^{\frac{1}{2}}\Big)\\
&= n\frac{\sum_{k=0}^{n}\frac{\lambda_n^k}{k!}}{\sum_{k=0}^{n-1}\frac{\lambda_n^k}{k!}} - \lambda_n\\
&> 0.
\end{aligned}$$

If $(\sqrt{n}-\sqrt{\lambda_n})^2 \geq 1$, then

$$\begin{aligned}
\Psi_n(1) &= \frac{\sum_{k=0}^{n}\binom{n}{k}\lambda_n^k\prod_{i=1}^{n-k}(i-1)}{\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{i=1}^{n-1-k}(i-1)} - \frac{1}{2}\Big(\lambda_n + n - 1 - \big((\lambda_n+n-1)^2 - 4\lambda_n n\big)^{\frac{1}{2}}\Big)\\
&= \frac{\lambda_n^n}{\lambda_n^{n-1}} - \frac{1}{2}\Big(\lambda_n + n - 1 - \big((\lambda_n+n-1)^2 - 4\lambda_n n\big)^{\frac{1}{2}}\Big)\\
&= \frac{1}{2}\Big(\lambda_n - n + 1 + \big((\lambda_n-n+1)^2 - 4\lambda_n\big)^{\frac{1}{2}}\Big)\\
&< \frac{1}{2}\big(\lambda_n - n + 1 + |\lambda_n - n + 1|\big)\\
&= 0 \quad \text{since} \;\; \lambda_n - n + 1 < 0.
\end{aligned}$$

We have

$$\begin{aligned}
\sigma_n(0) &= \sum_{k=0}^{n}\binom{n}{k}\lambda_n^k\prod_{i=1}^{n-k} i - \sqrt{\lambda_n n}\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{i=1}^{n-1-k} i\\
&= (n-1)!\big(n\sum_{k=0}^{n}\frac{\lambda_n^k}{k!} - \sqrt{\lambda_n n}\sum_{k=0}^{n-1}\frac{\lambda_n^k}{k!}\big)\\
&\geq (n-1)!\sum_{k=0}^{n}\frac{\lambda_n^k}{k!}(n - \sqrt{\lambda_n n})\\
&> 0,
\end{aligned}$$

and

$$\begin{aligned}
\sigma_n(1) &= \sum_{k=0}^{n}\binom{n}{k}\lambda_n^k\prod_{i=1}^{n-k}(i-1) - \sqrt{\lambda_n n}\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{i=1}^{n-1-k}(i-1)\\
&= \lambda_n^n - \sqrt{\lambda_n n}\lambda_n^{n-1}\\
&< 0.
\end{aligned}$$

We first prove assertion i. Since $(\sqrt{n}-\sqrt{\lambda_n})^2 < 1$, we have by (8) that $\min\big((\sqrt{n}-\sqrt{\lambda_n})^2, Z(f_{n,n-1})\big) = (\sqrt{n}-\sqrt{\lambda_n})^2$, and $f_{n,n-1}(x) > 0$ on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$. Thus from definitions (and dividing through by $f_{n,n-1}(y)$), we find that $\sigma_n(y)$ is the same sign as $z_n(y) - \sqrt{\lambda_n n}$, and that $\psi_n(y)$ is the same sign as $\Psi_n(y)$, on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$. Also, by Lemma 1.ii, $z_n(y)$ is strictly positive, continuous, and strictly decreasing on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$. Thus by the Intermediate Value Theorem, $\sigma_n(y)$ has a zero on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big)$ iff $z_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) - \sqrt{\lambda_n n} < 0$. By definitions and (10), we thus have that $Z(\sigma_n) < (\sqrt{n}-\sqrt{\lambda_n})^2$ iff $\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) < 0$. Since by assumption

$\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) < 0$, it will be the case that $Z(\sigma_n) < (\sqrt{n}-\sqrt{\lambda_n})^2$. Furthermore, it follows from the fact that $\Psi_n(0) > 0$ and the continuity and monotonicity of $\Psi_n(y)$ guaranteed by Lemma 1.iv that $Z(\Psi_n)$ will be the unique zero of Ψ_n in the interval $\big(0, (\sqrt{n}-\sqrt{\lambda_n})^2\big)$. Since $\psi_n(y)$ is the same sign as $\Psi_n(y)$ on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$, we will also have that $Z(\psi_n) = Z(\Psi_n)$. This, combined with Theorem 8, completes the proof of assertion i.

We now prove assertion ii. Since $(\sqrt{n}-\sqrt{\lambda_n})^2 < 1$, we have by (8) that $\min\big((\sqrt{n}-\sqrt{\lambda_n})^2, Z(f_{n,n-1})\big) = (\sqrt{n}-\sqrt{\lambda_n})^2$, and $f_{n,n-1}(x) > 0$ on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$. Thus from definitions (and dividing through by $f_{n,n-1}(y)$), we find that $\sigma_n(y)$ is the same sign as $z_n(y) - \sqrt{\lambda_n n}$, and that $\psi_n(y)$ is the same sign as $\Psi_n(y)$, on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$. Also, by Lemma 1.ii, $z_n(y)$ is strictly positive, continuous, and strictly decreasing on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big]$. Thus by the Intermediate Value Theorem, $\sigma_n(y)$ has a zero on $\big(-\infty, (\sqrt{n}-\sqrt{\lambda_n})^2\big)$ iff $z_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) - \sqrt{\lambda_n n} < 0$. By definitions and (10), we thus have that $Z(\sigma_n) < (\sqrt{n}-\sqrt{\lambda_n})^2$ iff $\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) < 0$. Since by assumption $\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) \geq 0$, it will be the case that $Z(\sigma_n) \geq (\sqrt{n}-\sqrt{\lambda_n})^2$. This, combined with Theorem 8, completes the proof of assertion ii.

We now prove assertion iii. By assumption, $(\sqrt{n}-\sqrt{\lambda_n})^2 \geq 1$. Thus since $Z(f_{n,n-1}) > 1$ by (8), we have that $f_{n,n-1}(x) > 0$ on $(-\infty, 1]$. Thus from definitions (and dividing through by $f_{n,n-1}(y)$), we find that $\psi_n(y)$ is the same sign as $\Psi_n(y)$ on $(-\infty, 1]$. Since $\sigma_n(y)$ is a polynomial and thus continuous, by the Intermediate Value Theorem and the fact that $\sigma_n(0) > 0$ and $\sigma_n(1) < 0$, we have $Z(\sigma_n) < 1$. Thus $Z(\sigma_n) < (\sqrt{n}-\sqrt{\lambda_n})^2$. By the continuity and monotonicity of $\Psi_n(y)$ guaranteed by Lemma 1.iv and the fact that $\Psi_n(0) > 0$, $\Psi_n(1) < 0$, we will have that $Z(\Psi_n)$ is the unique zero of Ψ_n in the interval $(0, 1)$. Furthermore, since $\psi_n(y)$ is the same sign as $\Psi_n(y)$ on $(-\infty, 1]$, we will have that $Z(\psi_n) = Z(\Psi_n)$. This, combined with Theorem 8, completing the proof of assertion iii. $\square$

We now prove some simple asymptotic bounds for use in later arguments.

Lemma 4. $\frac{B}{2} \leq \sqrt{n}-\sqrt{\lambda_n} \leq \frac{B}{2} + \frac{B^2}{4n^{\frac{1}{2}}-2B}$, *and* $\lim_{n\to\infty}(\sqrt{n}-\sqrt{\lambda_n}) = \frac{B}{2}$.

Proof.

$$\begin{aligned}\sqrt{n}-\sqrt{\lambda_n} &= \frac{(\sqrt{n}-\sqrt{\lambda_n})(\sqrt{n}+\sqrt{\lambda_n})}{\sqrt{n}+\sqrt{\lambda_n}} \\ &= \frac{B}{1+(\frac{\lambda_n}{n})^{\frac{1}{2}}} \\ &= \frac{B}{1+(1-Bn^{-\frac{1}{2}})^{\frac{1}{2}}}. \qquad (16)\end{aligned}$$

Note that

$$1 - Bn^{-\frac{1}{2}} \leq (1-Bn^{-\frac{1}{2}})^{\frac{1}{2}} \leq 1.$$

Combining with (16) we find that $\sqrt{n}-\sqrt{\lambda_n}$ is at least $\frac{B}{2}$, and at most

$$\frac{B}{2-Bn^{-\frac{1}{2}}} = \frac{B}{2} + \frac{B^2}{4n^{\frac{1}{2}}-2B}. \qquad (17)$$

The lemma follows letting $n \to \infty$ in (17). $\square$

Let $q_n \overset{\Delta}{=} 1 + n^{-\frac{1}{4}}$.

LEMMA 5. *For each $B > 0$, there exists N_B s.t. $n \geq N_B$ implies $g_n(k)\frac{n!}{\lambda_n^n} \leq q_n e^{B^2}$ for all k.*

PROOF. Since $\frac{\lambda_n}{n} < 1$, we have that $g_n(k) \leq \frac{\lambda_n^{\lfloor \lambda_n \rfloor}}{\lfloor \lambda_n \rfloor !}$ for all $k \geq 0$. Thus

$$\begin{aligned}
g_n(k)\frac{n!}{\lambda_n^n} &\leq \frac{\lambda_n^{\lfloor \lambda_n \rfloor}}{\lfloor \lambda_n \rfloor !}\frac{n!}{\lambda_n^n} \\
&\leq (\frac{n}{\lambda_n})^{n-\lfloor \lambda_n \rfloor} \\
&\leq (1+\frac{B}{n^{\frac{1}{2}}-B})^{Bn^{\frac{1}{2}}+1} \\
&\leq \exp\big(\frac{B^2 n^{\frac{1}{2}}}{n^{\frac{1}{2}}-B}+\frac{B}{n^{\frac{1}{2}}-B}\big) \\
&= \exp\big(B^2+\frac{B^3}{n^{\frac{1}{2}}-B}+\frac{B}{n^{\frac{1}{2}}-B}\big) \\
&= e^{B^2}\exp\big(O(n^{-\frac{1}{2}})\big) \\
&\leq q_n e^{B^2} \text{ for all sufficiently large n by a simple Taylor series expansion.}
\end{aligned}$$

□

LEMMA 6. *For all $0 < x < 1$, $T \geq 3$, $n \geq T+1$, and $k \in [0, n-T-1]$,*

$$e^{-2T^{-1}}(n-k)^{-x}T^x \leq \prod_{j=T+1}^{n-k}(1-\frac{x}{j}) \leq e^{2T^{-1}}(n-k)^{-x}T^x. \tag{18}$$

PROOF. Since $0 < x < 1$, by a simple Taylor series expansion we have that for $j \geq 3$, $1 \leq \frac{e^{-\frac{x}{j}}}{1-\frac{x}{j}} \leq 1+\frac{1}{j^2}$. Thus for $j \geq T+1$,

$$\begin{aligned}
\prod_{j=T+1}^{n-k}\frac{e^{-\frac{x}{j}}}{(1-\frac{x}{j})} &\leq \prod_{j=T+1}^{n-k}(1+\frac{1}{j^2}) \\
&\leq \prod_{j=T+1}^{\infty} e^{\frac{1}{j^2}} \\
&= e^{\sum_{j=T+1}^{\infty}\frac{1}{j^2}} \\
&\leq e^{\int_T^{\infty}\frac{1}{x^2}dx} \\
&= e^{T^{-1}},
\end{aligned}$$

and

$$e^{-T^{-1}}\prod_{j=T+1}^{n-k}e^{-\frac{x}{j}} \leq \prod_{j=T+1}^{n-k}(1-\frac{x}{j}) \leq \prod_{j=T+1}^{n-k}e^{-\frac{x}{j}}. \tag{19}$$

From [45], we have that for all $k \geq 1$, $\log(k)+\zeta \leq H_k \leq \frac{1}{2k}+\log(k)+\zeta$, and thus $\log(\frac{n-k}{T})-\frac{1}{2T} \leq H_{n-k}-H_T \leq \log(\frac{n-k}{T})+\frac{1}{2(n-k)}$. Furthermore, since by assumption $n-k > T$, we have

$$\log(\frac{n-k}{T})-\frac{1}{2T} \leq H_{n-k}-H_T \leq \log(\frac{n-k}{T})+\frac{1}{2T}. \tag{20}$$

Combining (19) and (20) yields

$$e^{-T^{-1}}e^{-\frac{x}{2T}}(n-k)^{-x}T^{x} \leq \prod_{j=T+1}^{n-k}(1-\frac{x}{j}) \leq e^{\frac{x}{2T}}(n-k)^{-x}T^{x}. \tag{21}$$

Since $0 < x < 1$ we have $0 < \frac{x}{2T} < \frac{1}{2T}$, and the lemma thus follows from (21). $\square$

We now state some additional properties of the parabolic cylinder functions. $D_x(z)$ is an entire function of z for fixed x ([15], Chapter 8, Section 8.2) and an entire function of x for fixed z [9]. It is stated in [21] that for all $x, z \in \mathbb{R}$,

$$D_{x+1}(z) - zD_x(z) + xD_{x-1}(z) = 0; \tag{22}$$

$$\frac{d}{dz}D_x(z) + \frac{1}{2}zD_x(z) - xD_{x-1}(z) = 0. \tag{23}$$

4. Asymptotics of $\Psi_n(x)$. In this section we derive the asymptotics of $\Psi_n(x)$ and $\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big)$. In particular, we prove

Theorem 10. *For all $B > 0$ and $x \in (0,1), x \leq \frac{B^2}{4}$, $\lim_{n\to\infty} \frac{\Psi_n(x)}{\sqrt{\lambda_n}} = \Psi_\infty(x)$.*

We also prove

Corollary 2. *For $0 < B < 2$, $\lim_{n\to\infty} \lambda_n^{-\frac{1}{2}}\Psi_n\big((\sqrt{n}-\sqrt{\lambda_n})^2\big) = \zeta(B)$.*

We proceed by separately analyzing the asymptotics of $\frac{a_n(x)-\lambda_n}{\sqrt{\lambda_n}}$ and $\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}$, and then use the fact that by definition $\frac{\Psi_n(x)}{\sqrt{\lambda_n}} = \frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} - \frac{a_n(x)-\lambda_n}{\sqrt{\lambda_n}}$.

We now analyze the asymptotics of $a_n(x)$.

Lemma 7. *For $0 \leq x \leq \frac{B^2}{4}$, $\lim_{n\to\infty} \frac{a_n(x)-\lambda_n}{\sqrt{\lambda_n}} = a_\infty(x)$.*

Proof. From (9) and Lemma 4,

$$\frac{a_n(x)-\lambda_n}{\sqrt{\lambda_n}} = \frac{B\sqrt{n} - x - \big((\sqrt{n}+\sqrt{\lambda_n})^2 - x\big)^{\frac{1}{2}}\big((\sqrt{n}-\sqrt{\lambda_n})^2 - x\big)^{\frac{1}{2}}}{2\sqrt{\lambda_n}}. \tag{24}$$

The lemma then follows from the fact that $\lim_{n\to\infty}\frac{B\sqrt{n}-x}{2\sqrt{\lambda_n}} = \frac{B}{2}$, $\lim_{n\to\infty}\frac{\big((\sqrt{n}+\sqrt{\lambda_n})^2-x\big)^{\frac{1}{2}}}{2\sqrt{\lambda_n}} = 1$, and Lemma 4. $\square$

We now analyze the asymptotics of $z_n(x)$, proving that

Proposition 2. *For $x \in \big(0,1\big)$, $\lim_{n\to\infty} \frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} = z_\infty(x)$.*

By (8), $f_{n,n-1}(x) > 0$ on $(0,1)$, and thus

$$
\begin{aligned}
\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} &= \frac{\frac{\sum_{k=0}^{n}\binom{n}{k}\lambda_n^k\prod_{j=1}^{n-k}(j-x)}{\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{j=1}^{n-1-k}(j-x)}-\lambda_n}{\sqrt{\lambda_n}} \\
&= \frac{\sum_{k=0}^{n}\binom{n}{k}\lambda_n^k\prod_{j=1}^{n-k}(j-x)-\lambda_n\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{j=1}^{n-1-k}(j-x)}{\sqrt{\lambda_n}\sum_{k=0}^{n-1}\binom{n-1}{k}\lambda_n^k\prod_{j=1}^{n-1-k}(j-x)}. \qquad (25)
\end{aligned}
$$

The remainder of the proof of Proposition 2 proceeds by a series of lemmas.

Lemma 8.

$$
\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} = \sqrt{\lambda_n}\,\frac{\sum_{k=0}^{n}(n-k)\prod_{j=1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}}{\sum_{k=0}^{n}k\prod_{j=1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}}. \qquad (26)
$$

Proof. We begin by analyzing the numerator of (25), which is equivalent to

$$
\begin{aligned}
&\prod_{j=1}^{n}(j-x)+\sum_{k=1}^{n}\binom{n}{k}\lambda_n^k\prod_{j=1}^{n-k}(j-x)-\sum_{k=0}^{n-1}\binom{n-1}{(k+1)-1}\lambda_n^{k+1}\prod_{j=1}^{n-(k+1)}(j-x) \\
&= \prod_{j=1}^{n}(j-x)+\sum_{k=1}^{n}\binom{n}{k}\lambda_n^k\prod_{j=1}^{n-k}(j-x)-\sum_{k=1}^{n}\binom{n-1}{k-1}\lambda_n^k\prod_{j=1}^{n-k}(j-x) \\
&= \prod_{j=1}^{n}(j-x)+\sum_{k=1}^{n}(\binom{n}{k}-\binom{n-1}{k-1})\lambda_n^k\prod_{j=1}^{n-k}(j-x) \\
&= \prod_{j=1}^{n}(j-x)+\sum_{k=1}^{n}\frac{(n-1)!}{(n-k)!k!}(n-k)\lambda_n^k\prod_{j=1}^{n-k}(j-x) \\
&= \prod_{j=1}^{n}(j-x)+(n-1)!\sum_{k=1}^{n}(n-k)\frac{\prod_{j=1}^{n-k}(j-x)}{(n-k)!}\frac{\lambda_n^k}{k!} \\
&= \prod_{j=1}^{n}(j-x)+(n-1)!\sum_{k=1}^{n}(n-k)\prod_{j=1}^{n-k}(1-\frac{x}{j})\ \frac{\lambda_n^k}{k!} \\
&= (n-1)!\big(n\prod_{j=1}^{n}(1-\frac{x}{j})+\sum_{k=1}^{n}(n-k)\prod_{j=1}^{n-k}(1-\frac{x}{j})\ \frac{\lambda_n^k}{k!}\big) \\
&= (n-1)!\sum_{k=0}^{n}(n-k)\prod_{j=1}^{n-k}(1-\frac{x}{j})\ \frac{\lambda_n^k}{k!}. \qquad (27)
\end{aligned}
$$

We now analyze the denominator of (25), which is equivalent to

$$
\begin{aligned}
\lambda_n^{-\frac{1}{2}}\sum_{k=0}^{n-1}\binom{n-1}{(k+1)-1}\lambda_n^{k+1}\prod_{j=1}^{n-(k+1)}(j-x) &= \lambda_n^{-\frac{1}{2}}\sum_{k=1}^{n}\frac{k}{n}\binom{n}{k}\lambda_n^k\prod_{j=1}^{n-k}(j-x) \\
&= \lambda_n^{-\frac{1}{2}}(n-1)!\sum_{k=0}^{n}k\prod_{j=1}^{n-k}(1-\frac{x}{j})\ \frac{\lambda_n^k}{k!}. \qquad (28)
\end{aligned}
$$

Combining (27) and (28), and normalizing both by $\frac{e^{-\lambda_n}}{(n-1)!}$, yields the lemma. □

Note that by a simple application of Stirling's Inequality, we find that $\lim_{n\to\infty}\sqrt{n}e^{-\lambda_n}\frac{\lambda_n^{\lfloor\lambda_n\rfloor}}{\lfloor\lambda_n\rfloor!} = (2\sqrt{\pi})^{-1}$. Thus we may fix a positive integer $T \geq 3$, and assume that n is sufficiently large to ensure that $\frac{n}{\lambda_n} \leq 2$, $\lceil n - T^{-1}\sqrt{n}\rceil + 1 \leq n-(T+1)$, $\frac{1}{\sqrt{\lambda_n}} \leq \frac{1}{2T}$, and $e^{-\lambda_n}\frac{\lambda_n^{\lfloor\lambda_n\rfloor}}{\lfloor\lambda_n\rfloor!} \leq n^{-\frac{1}{2}}$.

Lemma 9. *The numerator of (26) (ignoring the $\sqrt{\lambda_n}$ prefactor) is:*

(i) bounded from below by

$$\prod_{j=1}^{T}(1-\frac{x}{j})\sum_{k=0}^{n-(T+1)}(n-k)\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\; e^{-\lambda_n}\frac{\lambda_n^k}{k!}; \tag{29}$$

(ii) bounded from above by

$$\prod_{j=1}^{T}(1-\frac{x}{j})(\sum_{k=0}^{n-(T+1)}(n-k)\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\; e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\frac{(T+1)^2}{\prod_{j=1}^{T}(1-\frac{x}{j})\sqrt{n}}). \tag{30}$$

Proof. Note that the numerator of (26) is equivalent to

$$\prod_{j=1}^{T}(1-\frac{x}{j})\Bigg(\sum_{k=0}^{n-(T+1)}(n-k)\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\; e^{-\lambda_n}\frac{\lambda_n^k}{k!}+(\prod_{j=1}^{T}(1-\frac{x}{j}))^{-1}\sum_{k=n-T}^{n}(n-k)\prod_{j=1}^{n-k}(1-\frac{x}{j})\; e^{-\lambda_n}\frac{\lambda_n^k}{k!}\Bigg). \tag{31}$$

The second summand in (31) is non-negative, yielding a trivial lower bound. We now obtain an upper bound on the same expression. Since the mode of a Poisson r.v. with mean λ_n is $\lfloor\lambda_n\rfloor$, and by assumption $e^{-\lambda_n}\frac{\lambda_n^{\lfloor\lambda_n\rfloor}}{\lfloor\lambda_n\rfloor!} \leq n^{-\frac{1}{2}}$, we have that for all $k \geq 0$,

$$e^{-\lambda_n}\frac{\lambda_n^k}{k!} \leq e^{-\lambda_n}\frac{\lambda_n^{\lfloor\lambda_n\rfloor}}{\lfloor\lambda_n\rfloor!} \leq n^{-\frac{1}{2}}. \tag{32}$$

We now use the fact that $n-k \leq T+1$ for $k \geq n-T$, $1-\frac{x}{j} \leq 1$, and (32) to conclude that the second summand in (31) is at most

$$\frac{(T+1)^2}{\prod_{j=1}^{T}(1-\frac{x}{j})\sqrt{n}}. \tag{33}$$

Combining our lower and upper bounds on (31) completes the proof of the lemma. □

Lemma 10. *The denominator of (26) is:*

(i) bounded from below by

$$\prod_{j=1}^{T}(1-\frac{x}{j})\sum_{k=0}^{n-(T+1)}k\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\; e^{-\lambda_n}\frac{\lambda_n^k}{k!}; \tag{34}$$

(ii) bounded from above by

$$\prod_{j=1}^{T}(1-\frac{x}{j})(\sum_{k=0}^{n-(T+1)} k \prod_{j=T+1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\frac{(T+1)\sqrt{n}}{\prod_{j=1}^{T}(1-\frac{x}{j})}). \tag{35}$$

PROOF. Note that the denominator of (26) is equivalent to

$$\prod_{j=1}^{T}(1-\frac{x}{j})(\sum_{k=0}^{n-(T+1)} k \prod_{j=T+1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+(\prod_{j=1}^{T}(1-\frac{x}{j}))^{-1}\sum_{k=n-T}^{n} k\prod_{j=1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}). \tag{36}$$

We now bound the second summand in (36). Note that it is non-negative, yielding a trivial lower bound. Also, it is at most

$$\begin{aligned}\frac{T+1}{\prod_{j=1}^{T}(1-\frac{x}{j})}\max_{n-T\leq k\leq n} k\prod_{j=1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!} &\leq \frac{T+1}{\prod_{j=1}^{T}(1-\frac{x}{j})}\ n\ \frac{1}{\sqrt{n}} \quad \text{by (32)}\\ &\leq \frac{(T+1)\sqrt{n}}{\prod_{j=1}^{T}(1-\frac{x}{j})}.\end{aligned} \tag{37}$$

Combining our lower and upper bounds on (36) completes the proof of the lemma. ☐

Let $C_{T,x}\triangleq\max(\frac{2(T+1)T^{-x}}{\prod_{j=1}^{T}(1-\frac{x}{j})},\frac{(T+1)^2T^{-x}}{\prod_{j=1}^{T}(1-\frac{x}{j})})$.

LEMMA 11.

(i)

$$\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}\geq\frac{e^{-2T^{-1}}\sum_{k=0}^{n-(T+1)}(n-k)^{1-x}e^{-\lambda_n}\frac{\lambda_n^k}{k!}}{e^{2T^{-1}}\sum_{k=0}^{n-(T+1)}\frac{k}{\sqrt{\lambda_n}}(n-k)^{-x}\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+C_{T,x}}; \tag{38}$$

(ii)

$$\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}\leq\frac{e^{2T^{-1}}\sum_{k=0}^{n-(T+1)}(n-k)^{1-x}\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+C_{T,x}n^{-\frac{1}{2}}}{e^{-2T^{-1}}\sum_{k=0}^{n-(T+1)}\frac{k}{\sqrt{\lambda_n}}(n-k)^{-x}\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}}. \tag{39}$$

PROOF. Combining Lemma 9 and Lemma 10 we obtain

$$\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}\geq\sqrt{\lambda_n}\frac{\sum_{k=0}^{n-(T+1)}(n-k)\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}}{\sum_{k=0}^{n-(T+1)}k\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\frac{(T+1)\sqrt{n}}{\prod_{j=1}^{T}(1-\frac{x}{j})}}, \tag{40}$$

and

$$\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}\leq\sqrt{\lambda_n}\frac{\sum_{k=0}^{n-(T+1)}(n-k)\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\frac{(T+1)^2}{\prod_{j=1}^{T}(1-\frac{x}{j})\sqrt{n}}}{\sum_{k=0}^{n-(T+1)}k\prod_{j=T+1}^{n-k}(1-\frac{x}{j})\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}}. \tag{41}$$

Combining (40) and (41) with Lemma 6 and the assumptions on n, we find that

$$\begin{aligned}\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} &\geq \sqrt{\lambda_n}\frac{\sum_{k=0}^{n-(T+1)}(n-k)\big(e^{-2T^{-1}}(n-k)^{-x}T^{x}\big)e^{-\lambda_n}\frac{\lambda_n^k}{k!}}{\sum_{k=0}^{n-(T+1)}k\big(e^{2T^{-1}}(n-k)^{-x}T^{x}\big)\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\frac{(T+1)\sqrt{n}}{\prod_{j=1}^{T}(1-\frac{x}{j})}} \\ &= \frac{e^{-2T^{-1}}\sum_{k=0}^{n-(T+1)}(n-k)^{1-x}e^{-\lambda_n}\frac{\lambda_n^k}{k!}}{e^{2T^{-1}}\sum_{k=0}^{n-(T+1)}\frac{k}{\sqrt{\lambda_n}}(n-k)^{-x}\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\sqrt{\frac{n}{\lambda_n}}\frac{(T+1)T^{-x}}{\prod_{j=1}^{T}(1-\frac{x}{j})}},\end{aligned}$$

from which (38) follows since $\frac{n}{\lambda_n}\leq 2$. (39) follows from a nearly identical argument, proving the lemma. ☐

LEMMA 12.

(i)

$$\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} \geq \frac{e^{-2T^{-1}}\lambda_n^{\frac{x-1}{2}}\sum_{k=0}^{n-(T+1)}(n-k)^{1-x}e^{-\lambda_n}\frac{\lambda_n^k}{k!}}{e^{2T^{-1}}\lambda_n^{\frac{x-1}{2}}\sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\frac{2e^{2T^{-1}}T^{-(1-x)}(\frac{n}{\lambda_n})^{\frac{1-x}{2}}}{1-x}+\lambda_n^{\frac{x-1}{2}}C_{T,x}}; \tag{42}$$

(ii)

$$\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} \leq \frac{e^{2T^{-1}}\lambda_n^{\frac{x-1}{2}}\sum_{k=0}^{n-(T+1)}(n-k)^{1-x}e^{-\lambda_n}\frac{\lambda_n^k}{k!}+\lambda_n^{\frac{x-1}{2}}C_{T,x}n^{-\frac{1}{2}}}{e^{-2T^{-1}}\lambda_n^{\frac{x-1}{2}}\sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}}. \tag{43}$$

PROOF. We first prove i. The term $\sum_{k=0}^{n-(T+1)}\frac{k}{\sqrt{\lambda_n}}(n-k)^{-x}\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}$ appearing in the denominator of (38) is at most

$$\sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\max_{\lceil n-T^{-1}\sqrt{n}\rceil+1\leq k\leq n}\left(\frac{k}{\sqrt{\lambda_n}}\ e^{-\lambda_n}\frac{\lambda_n^k}{k!}\right)\sum_{k=\lceil n-T^{-1}\sqrt{n}\rceil+1}^{n}(n-k)^{-x}$$

$$\begin{aligned}&\leq \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\frac{n}{\sqrt{\lambda_n}}n^{-\frac{1}{2}}\sum_{k=\lceil n-T^{-1}\sqrt{n}\rceil+1}^{n}(n-k)^{-x} \quad \text{by (32)} \\ &\leq \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\sqrt{\frac{n}{\lambda_n}}\sum_{j=T+1}^{\lceil T^{-1}\sqrt{n}\rceil-1}j^{-x} \\ &\leq \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\sqrt{\frac{n}{\lambda_n}}\int_0^{T^{-1}\sqrt{n}}y^{-x}dy \\ &\leq \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\sqrt{\frac{n}{\lambda_n}}n^{\frac{1-x}{2}}\frac{T^{-(1-x)}}{1-x} \\ &\leq \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil}\frac{k(n-k)^{-x}}{\sqrt{\lambda_n}}\frac{e^{-\lambda_n}\lambda_n^k}{k!}+\frac{2T^{-(1-x)}n^{\frac{1-x}{2}}}{1-x} \quad \text{since } \frac{n}{\lambda_n}\leq 2. \qquad (44)\end{aligned}$$

We may thus upper-bound the denominator of (38) using (44), and then multiply both numerator and denominator by $\lambda_n^{\frac{x-1}{2}}$, to prove i.

We now prove ii. By non-negativity the denominator of (39) is at least $e^{-2T^{-1}} \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil} \frac{k(n-k)^{-x}}{\sqrt{\lambda_n}} \frac{e^{-\lambda_n}\lambda_n^k}{k!}$, and thus ii. follows from (39) and multiplying both numerator and denominator by $\lambda_n^{\frac{x-1}{2}}$. $\square$

We now relate the summations appearing in (42) and (43) to the expectations of certain functions of a poisson r.v., and then show that these expectations converge to certain integrals as $n \to \infty$. Let X_n denote a Poisson r.v. with mean λ_n, $Z_n \triangleq \frac{X_n - \lambda_n}{\sqrt{\lambda_n}}$, and $I(G)$ denote the indicator of the event G. Thus

$$X_n = \sqrt{\lambda_n} Z_n + \lambda_n, \tag{45}$$

and

$$\begin{aligned} n - X_n &= n - (\sqrt{\lambda_n} Z_n + \lambda_n) \\ &= B\sqrt{n} - \sqrt{\lambda_n} Z_n. \end{aligned} \tag{46}$$

Furthermore, let

$$Y_{1,n} \triangleq (B\sqrt{\frac{n}{\lambda_n}} - Z_n)^{1-x} I(Z_n \leq B\sqrt{\frac{n}{\lambda_n}} - \frac{T+1}{\sqrt{\lambda_n}}), \tag{47}$$

$$Y_{2,n} \triangleq Z_n (B\sqrt{\frac{n}{\lambda_n}} - Z_n)^{-x} I(Z_n \leq (B - T^{-1})\sqrt{\frac{n}{\lambda_n}} + \frac{\lceil n - T^{-1}\sqrt{n}\rceil - (n - T^{-1}\sqrt{n})}{\sqrt{\lambda_n}}), \tag{48}$$

and

$$Y_{3,n} \triangleq (B\sqrt{\frac{n}{\lambda_n}} - Z_n)^{-x} I(Z_n \leq (B - T^{-1})\sqrt{\frac{n}{\lambda_n}} + \frac{\lceil n - T^{-1}\sqrt{n}\rceil - (n - T^{-1}\sqrt{n})}{\sqrt{\lambda_n}}). \tag{49}$$

We then have that

LEMMA 13. *If* $\lim_{n\to\infty} E[Y_{1,n}]$, $\lim_{n\to\infty} E[Y_{2,n}]$, *and* $\lim_{n\to\infty} E[Y_{3,n}]$ *all exist, and are finite and strictly positive, then*

(i)

$$\liminf_{n\to\infty} \frac{z_n(x) - \lambda_n}{\sqrt{\lambda_n}} \geq e^{-4T^{-1}} \frac{\lim_{n\to\infty} E[Y_{1,n}]}{\lim_{n\to\infty} E[Y_{3,n}] + \frac{2T^{-(1-x)}}{1-x}}; \tag{50}$$

(ii)

$$\limsup_{n\to\infty} \frac{z_n(x) - \lambda_n}{\sqrt{\lambda_n}} \leq e^{4T^{-1}} \frac{\lim_{n\to\infty} E[Y_{1,n}]}{\lim_{n\to\infty} E[Y_{3,n}]}. \tag{51}$$

PROOF. We begin by expressing the summation appearing in the numerator of (42) and (43) as an expectation.

$$\begin{aligned} \lambda_n^{\frac{x-1}{2}} \sum_{k=0}^{n-(T+1)} (n-k)^{1-x} e^{-\lambda_n} \frac{\lambda_n^k}{k!} &= \lambda_n^{\frac{x-1}{2}} E[(n - X_n)^{1-x} I\big(X_n \leq n - (T+1)\big)] \\ &= \lambda_n^{\frac{x-1}{2}} E[(B\sqrt{n} - \sqrt{\lambda_n} Z_n)^{1-x} I\big(\sqrt{\lambda_n} Z_n + \lambda_n \leq n - (T+1)\big)] \quad \text{by (46)} \\ &= E[Y_{1,n}]. \end{aligned} \tag{52}$$

We now express the summation appearing in the denominator of (42) and (43) as the sum of two expectations.

$$\lambda_n^{\frac{x-1}{2}} \sum_{k=0}^{\lceil n-T^{-1}\sqrt{n}\rceil} \frac{k(n-k)^{-x}}{\sqrt{\lambda_n}} \frac{e^{-\lambda_n}\lambda_n^k}{k!}$$

$$\begin{aligned}
&= \lambda_n^{\frac{x-2}{2}} E[X_n(n-X_n)^{-x} I(X_n \leq \lceil n-T^{-1}\sqrt{n}\rceil)] \\
&= \lambda_n^{\frac{x-2}{2}} E[(\sqrt{\lambda_n}Z_n+\lambda_n)(B\sqrt{n}-\sqrt{\lambda_n}Z_n)^{-x} I(\sqrt{\lambda_n}Z_n+\lambda_n \leq \lceil n-T^{-1}\sqrt{n}\rceil)] \\
&\quad \text{by (45) and (46)} \\
&= \lambda_n^{\frac{x-1}{2}} E[(Z_n+\sqrt{\lambda_n})(B\sqrt{n}-\sqrt{\lambda_n}Z_n)^{-x} I(\sqrt{\lambda_n}Z_n+\lambda_n \leq \lceil n-T^{-1}\sqrt{n}\rceil)]. \qquad (53)
\end{aligned}$$

We now manipulate the indicator variable appearing in (53) to make it more amenable to asymptotic analysis. The indicator is

$$I(Z_n \leq \frac{\lceil n-T^{-1}\sqrt{n}\rceil - \lambda_n}{\sqrt{\lambda_n}})$$

$$\begin{aligned}
&= I(Z_n \leq \frac{\lceil n-T^{-1}\sqrt{n}\rceil - (n-B\sqrt{n})}{\sqrt{\lambda_n}}) \\
&= I(Z_n \leq \frac{\lceil n-T^{-1}\sqrt{n}\rceil - (n-T^{-1}\sqrt{n}) + (B-T^{-1})\sqrt{n}}{\sqrt{\lambda_n}}) \\
&= I(Z_n \leq (B-T^{-1})\sqrt{\frac{n}{\lambda_n}} + \frac{\lceil n-T^{-1}\sqrt{n}\rceil - (n-T^{-1}\sqrt{n})}{\sqrt{\lambda_n}}). \qquad (54)
\end{aligned}$$

Plugging (54) back into (53), we find that the summation appearing in the numerator of (42) and (43) is

$$\lambda_n^{-\frac{1}{2}} E[(Z_n+\sqrt{\lambda_n})(B\sqrt{\frac{n}{\lambda_n}}-Z_n)^{-x} I(Z_n \leq (B-T^{-1})\sqrt{\frac{n}{\lambda_n}} + \frac{\lceil n-T^{-1}\sqrt{n}\rceil - (n-T^{-1}\sqrt{n})}{\sqrt{\lambda_n}})]$$

(55)
$= \lambda_n^{-\frac{1}{2}} E[Y_{2,n}] + E[Y_{3,n}]$ by breaking up the $(Z_n+\sqrt{\lambda_n})$ term and applying linearity of expectations.

The lemma then follows by plugging (52) and (55) into (42) and (43), and observing that

$$\lim_{n\to\infty} \frac{2e^{2T^{-1}}T^{-(1-x)}(\frac{n}{\lambda_n})^{\frac{1-x}{2}}}{1-x} = \frac{2e^{2T^{-1}}T^{-(1-x)}}{1-x}, \quad \lim_{n\to\infty} \lambda_n^{\frac{x-1}{2}} C_{T,x} = 0, \text{ and } \lim_{n\to\infty} \lambda_n^{-\frac{1}{2}} E[Y_{2,n}] = 0.$$

□

We now explicitly compute the limiting values of all expressions appearing in (50) and (51), allowing us to compute $\lim_{n\to\infty} \frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}$.

Lemma 14.

$$\lim_{n\to\infty} \frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}} = \frac{\int_{-\infty}^{B} (B-z)^{1-x} e^{-\frac{z^2}{2}} dz}{\int_{-\infty}^{B} (B-z)^{-x} e^{-\frac{z^2}{2}} dz}. \qquad (56)$$

Proof. It is easily verified that $\{Y_{1,n}\},\{Y_{2,n}\},\{Y_{3,n}\}$ are all sequences of uniformly integrable r.v. Let $f_1(y) \triangleq (B-y)^{1-x}I(y \leq B), f_2(y) \triangleq y(B-y)^{-x}I(y \leq B-T^{-1}), f_3(y) \triangleq (B-y)^{-x}I(y \leq B-T^{-1})$. Let N denote a normal r.v. with mean 0 and variance 1. It may be easily verified that $\{Y_{1,n}\},\{Y_{2,n}\},\{Y_{3,n}\}$ converge weakly to $f_1(N), f_2(N), f_3(N)$ respectively, where all three convergences follow from the well-known convergence of the scaled Poisson r.v. Z_n to the Normal r.v. N. It thus follows from the uniform integrability of all three sequences that

$$\lim_{n\to\infty} E[Y_{1,n}] = E[f_1(N)] = (2\pi)^{-\frac{1}{2}}\int_{-\infty}^{B}(B-z)^{1-x}e^{-\frac{z^2}{2}}dz, \tag{57}$$

$$\lim_{n\to\infty} E[Y_{2,n}] = E[f_2(N)] = (2\pi)^{-\frac{1}{2}}\int_{-\infty}^{B-T^{-1}} z(B-z)^{-x}e^{-\frac{z^2}{2}}dz, \tag{58}$$

and

$$\lim_{n\to\infty} E[Y_{3,n}] = E[f_3(N)] = (2\pi)^{-\frac{1}{2}}\int_{-\infty}^{B-T^{-1}} (B-z)^{-x}e^{-\frac{z^2}{2}}dz. \tag{59}$$

The lemma follows from plugging the limits (57)-(59) into Lemma 13, realizing that T was general, and letting $T \to \infty$. □

We now relate the integrals appearing in (56) to the parabolic cylinder functions. In particular, we prove

Lemma 15. $\frac{\int_{-\infty}^{B}(B-z)^{1-x}e^{-\frac{z^2}{2}}dz}{\int_{-\infty}^{B}(B-z)^{-x}e^{-\frac{z^2}{2}}dz} = \frac{D_x(-B)}{D_{x-1}(-B)} + B.$

Proof. Note that the right-hand side of (56) is equivalent to

$$\begin{aligned}
\frac{\int_0^\infty t^{1-x}e^{-\frac{(B-t)^2}{2}}dt}{\int_0^\infty t^{-x}e^{-\frac{(B-t)^2}{2}}dt} &= \frac{e^{-\frac{B^2}{2}}\int_0^\infty t^{1-x}e^{-\frac{t^2}{2}+Bt}dt}{e^{-\frac{B^2}{2}}\int_0^\infty t^{-x}e^{-\frac{t^2}{2}+Bt}dt} \\
&= \frac{\int_0^\infty t^{-\left((x-2)+1\right)}e^{-\frac{t^2}{2}-(-B)t}dt}{\int_0^\infty t^{-\left((x-1)+1\right)}e^{-\frac{t^2}{2}+Bt}dt} \\
&= \frac{e^{-\frac{(-B)^2}{4}}\int_0^\infty t^{-\left((x-2)+1\right)}e^{-\frac{t^2}{2}-(-B)t}dt}{e^{-\frac{(-B)^2}{4}}\int_0^\infty t^{-\left((x-1)+1\right)}e^{-\frac{t^2}{2}-(-B)t}dt} \\
&= \frac{\Gamma\big(-(x-2)\big)D_{x-2}(-B)}{\Gamma\big(-(x-1)\big)D_{x-1}(-B)} \quad \text{by (2), since } \ x \in (0,1) \\
&= \frac{\Gamma(2-x)\frac{\left(D_x(-B)+BD_{x-1}(-B)\right)}{1-x}}{\Gamma(1-x)D_{x-1}(-B)} \quad \text{by (22)} \\
&= \frac{D_x(-B)}{D_{x-1}(-B)} + B \quad \text{since } \Gamma(2-x) = (1-x)\Gamma(1-x). \qquad (60)
\end{aligned}$$

□

We now complete the proofs of Proposition 2, Theorem 10, and Corollary 2.

Proof of Proposition 2. Proposition 2 follows by combining Lemma 14 and Lemma 15. $\square$

Proof of Theorem 10. Since by definition $\Psi_n(x) = z_n(x) - a_n(x)$, Theorem 10 follows immediately from Lemma 7 and Proposition 2. $\square$

Proof of Corollary 2. Suppose $0 < \epsilon < \min(\frac{B^2}{4}, 1 - \frac{B^2}{4})$. Then for all sufficiently large n, $0 < \frac{B^2}{4} - \epsilon < (\sqrt{n} - \sqrt{\lambda_n})^2 < \min(1-\epsilon, \frac{B^2}{4} + \epsilon)$ by Lemma 4, and $\frac{B^2}{4} < 1-\epsilon$. By Lemma 1.ii and (8), we have that $\frac{z_n(x)-\lambda_n}{\sqrt{\lambda_n}}$ is a continuous, strictly decreasing function of x on $(-\infty, (\sqrt{n} - \sqrt{\lambda_n})^2 + \epsilon]$. It follows that for all sufficiently large n, $\frac{z_n(\frac{B^2}{4}+\epsilon)-\lambda_n}{\sqrt{\lambda_n}} \leq \frac{z_n\left((\sqrt{n}-\sqrt{\lambda_n})^2\right)-\lambda_n}{\sqrt{\lambda_n}} \leq \frac{z_n(\frac{B^2}{4}-\epsilon)-\lambda_n}{\sqrt{\lambda_n}}$. Thus by Proposition 2, for all sufficiently small $\epsilon > 0$,

(61)
$$z_\infty(\frac{B^2}{4} + \epsilon) \leq \liminf_{n\to\infty} \frac{z_n\left((\sqrt{n} - \sqrt{\lambda_n})^2\right) - \lambda_n}{\sqrt{\lambda_n}} \leq \limsup_{n\to\infty} \frac{z_n\left((\sqrt{n} - \sqrt{\lambda_n})^2\right) - \lambda_n}{\sqrt{\lambda_n}} \leq z_\infty(\frac{B^2}{4} - \epsilon).$$

To proceed, we will now prove that $z_\infty(x)$ has certain continuity properties. In particular,

(62) $z_\infty(x)$ is a differentiable, non-increasing function of x on $(-\infty, 1]$.

Indeed, by (4), $D_{x-1}(-B) > 0$ for $x \leq 1$. The differentiability of $z_\infty(x)$ on $(-\infty, 1]$ then follows from the fact that $D_x(-B)$ is an entire function of x. That $z_\infty(x)$ is non-increasing follows from Lemma 1.ii, (8), and Proposition 2.

The continuity of $z_\infty(x)$ in a neighborhood of $\frac{B^2}{4}$ (guaranteed by (62) since $0 < \frac{B^2}{4} < 1$), along with (61), thus implies that $\lim_{n\to\infty} \frac{z_n\left((\sqrt{n}-\sqrt{\lambda_n})^2\right)-\lambda_n}{\sqrt{\lambda_n}} = z_\infty(\frac{B^2}{4})$. By (10), $\frac{a_n\left((\sqrt{n}-\sqrt{\lambda_n})^2\right)-\lambda_n}{\sqrt{\lambda_n}} = \sqrt{n} - \sqrt{\lambda_n}$. Thus by Lemma 4, $\lim_{n\to\infty} \frac{a_n\left((\sqrt{n}-\sqrt{\lambda_n})^2\right)-\lambda_n}{\sqrt{\lambda_n}} = \frac{B}{2}$, and the corollary follows since $\zeta(B) = \Psi_\infty(\frac{B^2}{4})$. $\square$

5. Asymptotics of $Z^+(\Psi_n)$.

In this section we derive the asymptotics of $Z^+(\Psi_n)$. In particular, we prove

Theorem 11. *If $B < 2$ and $\zeta(B) < 0$, or $B \geq 2$, then $\lim_{n\to\infty} Z^+(\Psi_n) = Z^+(\Psi_\infty)$.*

We begin by establishing some additional properties of $\Psi_\infty(x)$.

Lemma 16. *$\Psi_\infty(x)$ is a continuous function of x on $\left(-\infty, \min(1, \frac{B^2}{4})\right)$, left-continuous at $\min(1, \frac{B^2}{4})$, and a differentiable function of x on $\left(-\infty, \min(1, \frac{B^2}{4})\right)$, satisfying $\frac{d}{dx}\Psi_\infty(x) \leq -(B^2 - 4x)^{-1}$ for $x < \min(1, \frac{B^2}{4})$.*

Proof. That $a_\infty(x)$ is differentiable on $(-\infty, \frac{B^2}{4})$, left-continuous at $\frac{B^2}{4}$, and satisfies $\frac{d}{dx}a_\infty(x) = (B^2 - 4x)^{-1}$ for $x < \frac{B^2}{4}$, follows from elementary calculus. The lemma then follows from (62) and the fact that $\Psi_\infty(x) = z_\infty(x) - a_\infty(x)$. $\square$

We now prove some additional properties of $Z^+(\Psi_\infty)$.

Lemma 17.

(i) If $B < 2$ *and* $\zeta(B) < 0$*, or* $B \geq 2$*,*

  (a) $\Psi_\infty(x)$ *has a unique zero* $Z^+(\Psi_\infty) \in \big(0, \min(1, \frac{B^2}{4})\big)$*,*

  (b) $\Psi_\infty(x) > 0$ *on* $[0, Z^+(\Psi_\infty)\big)$*,*

  (c) $\Psi_\infty(x) < 0$ *on* $\big(Z^+(\Psi_\infty), \min(1, \frac{B^2}{4})]$.

(ii) If $B < 2$ *and* $\zeta(B) = 0$*,*

  (a) $\Psi_\infty(x) > 0$ *on* $[0, \min(1, \frac{B^2}{4})\big)$*,*

  (b) $Z^+(\Psi_\infty) = \frac{B^2}{4}$.

Proof. We first prove assertion i. By Lemma 16, $\Psi_\infty(x)$ is a strictly decreasing function of x on $[0, \min(1, \frac{B^2}{4})\big)$, continuous on $[0, \min(1, \frac{B^2}{4})\big)$, and left-continuous at $\min(1, \frac{B^2}{4})$. $\Psi_\infty(0) > 0$, since $\Psi_\infty(0) = \frac{D_0(-B)}{D_{-1}(-B)} + B$, and by (3), $\frac{D_0(-B)}{D_{-1}(-B)} > 0$.

Also, $\Psi_\infty(\min(1, \frac{B^2}{4})) < 0$. Indeed, if $B \geq 2$, $\min(1, \frac{B^2}{4}) = 1$. Furthermore, $\Psi_\infty(1) < 0$, since $\Psi_\infty(1) = \frac{D_1(-B)}{D_0(-B)} + \frac{1}{2}\big(B + (B^2-4)^{\frac{1}{2}}\big)$, and by (3),

$$\begin{aligned} \frac{D_1(-B)}{D_0(-B)} + \frac{1}{2}\big(B + (B^2-4)^{\frac{1}{2}}\big) &= -B + \frac{1}{2}\big(B + (B^2-4)^{\frac{1}{2}}\big) \\ &= \frac{1}{2}\big((B^2-4)^{\frac{1}{2}} - B\big) \\ &< 0. \end{aligned}$$

If $B < 2$, then $\min(1, \frac{B^2}{4}) = \frac{B^2}{4}$, and $\Psi_\infty(\frac{B^2}{4}) = \zeta(B) < 0$. Combining the cases $B \leq 2$ and $B > 2$ demonstrates that $\Psi_\infty(\min(1, \frac{B^2}{4})) < 0$.

Thus $\Psi_\infty(0) > 0$, and $\Psi_\infty(\min(1, \frac{B^2}{4})) < 0$. That $\Psi_\infty(x)$ has a unique zero $Z^+(\Psi_\infty) \in \big(0, \min(1, \frac{B^2}{4})\big)$, $\Psi_\infty(x) > 0$ on $[0, Z^+(\Psi_\infty)\big)$, and $\Psi_\infty(x) < 0$ on $\big(Z^+(\Psi_\infty), \min(1, \frac{B^2}{4})]$ then follows from the continuity and strict monotonicity of $\Psi_\infty(x)$ on $\big(-\infty, \min(1, \frac{B^2}{4})]$ guaranteed by Lemma 16.

We now prove assertion ii. The lemma follows from the fact that $\Psi_\infty(x)$ is strictly decreasing on $\big(-\infty, \min(1, \frac{B^2}{4})\big)$ and left-continuous at $\frac{B^2}{4}$ by Lemma 16, and $\zeta(B) = 0$. □

Proof of Theorem 11. We first show that for all sufficiently large n, $Z^+(\Psi_n)$ is the unique zero of Ψ_n in the interval $(0, \min(1, (\sqrt{n} - \sqrt{\lambda_n})^2)$. Indeed, if $B < 2$ and $\zeta(B) < 0$, then for all sufficiently large n, $\min(1, (\sqrt{n} - \sqrt{\lambda_n})^2)\big) = (\sqrt{n} - \sqrt{\lambda_n})^2$ by Lemma 4 and $\Psi_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) < 0$ by Corollary 2. It follows from Lemma 3.i that for all sufficiently large n, $Z^+(\Psi_n)$ is the unique zero of Ψ_n in the interval $\min(1, (\sqrt{n} - \sqrt{\lambda_n})^2)$. If instead $B \geq 2$, then for all sufficiently large n, $(\sqrt{n} - \sqrt{\lambda_n})^2 \geq 1$ and $\min(1, (\sqrt{n} - \sqrt{\lambda_n})^2)\big) = 1$ by Lemma 4. It follows from Lemma 3.iii that for all sufficiently large n, $Z^+(\Psi_n)$ is the unique zero of Ψ_n in the interval $(0, \min(1, (\sqrt{n} - \sqrt{\lambda_n})^2))$.

Now suppose for contradiction that $\liminf_{n\to\infty} Z^+(\Psi_n) < Z^+(\Psi_\infty)$. By Lemma 17.i, $Z^+(\Psi_\infty) \in \big(0, \min(1, \frac{B^2}{4})\big)$, and $\Psi_\infty(x) > 0$ on $[0, Z^+(\Psi_\infty)\big)$. It follows that there exists $\epsilon > 0$ such that $0 < \liminf_{n\to\infty} Z^+(\Psi_n) + \epsilon < \min(1, \frac{B^2}{4})$, and $\Psi_\infty(\liminf_{n\to\infty} Z^+(\Psi_n) + \epsilon) > 0$. Thus by Theorem 10, for all sufficiently large n, $\Psi_n(\liminf_{n\to\infty} Z^+(\Psi_n) + \epsilon) > 0$. By the monotonicity of $\Psi_n(x)$ (from Lemma 1.iv), it follows that for all sufficiently large n, $\Psi_n(x) > 0$ on $(-\infty, \liminf_{n\to\infty} Z^+(\Psi_n) + \epsilon)$. But by the definition of $\liminf$, there exists an infinite strictly increasing sequence of integers $\{n_i\}$

s.t. $Z^+(\Psi_{n_i}) < \liminf_{n\to\infty} Z^+(\Psi_n) + \epsilon$ for all i. Thus for all sufficiently large i, $\Psi_{n_i}(Z^+(\Psi_{n_i})) > 0$. This is a contradiction, since from definitions $\Psi_{n_i}(Z^+(\Psi_{n_i})) = 0$ for all i.

Alternatively, suppose for contradiction that $\limsup_{n\to\infty} Z^+(\Psi_n) > Z^+(\Psi_\infty)$. By Lemma 17.i, $Z^+(\Psi_\infty) \in \big(0, \min(1, \frac{B^2}{4})\big)$, and $\Psi_\infty(x) < 0$ on $\big(Z^+(\Psi_\infty), \min(1, \frac{B^2}{4})\big]$. It follows that there exists $\epsilon > 0$ such that $0 < \limsup_{n\to\infty} Z^+(\Psi_n) - \epsilon < \min(1, \frac{B^2}{4})$, and $\Psi_\infty(\limsup_{n\to\infty} Z^+(\Psi_n) - \epsilon) < 0$. Thus by Theorem 10, for all sufficiently large n, $\Psi_n(\limsup_{n\to\infty} Z^+(\Psi_n) - \epsilon) < 0$. By the monotonicity of $\Psi_n(x)$ (from Lemma 1.iv), it follows that for all sufficiently large n, $\Psi_n(x) < 0$ on $\big(\limsup_{n\to\infty} Z^+(\Psi_n) - \epsilon, \min(1, (\sqrt{n} - \sqrt{\lambda_n})^2)\big)$.

By the definition of $\limsup$, there exists an infinite strictly increasing sequence of integers $\{n_i\}$ s.t. $Z^+(\Psi_{n_i}) > \limsup_{n\to\infty} Z^+(\Psi_n) - \epsilon$ for all i. Combining the above, we find that for all sufficiently large i, $\Psi_{n_i}(Z^+(\Psi_{n_i})) < 0$. But this is a contradiction, since from definitions $\Psi_{n_i}(Z^+(\Psi_{n_i})) = 0$ for all i. We conclude that for the case $B < 2$ and $\zeta(B) < 0$, or $B \geq 2$, $\liminf_{n\to\infty} Z^+(\Psi_n) = \limsup_{n\to\infty} Z^+(\Psi_n) = Z^+(\Psi_\infty)$. ☐

6. The Zeros of $\zeta(B)$.

In this section we characterize the set of $B \in (0, 2)$ for which $\zeta(B) \leq 0$, which will allow us to use Theorem 11 effectively. In particular, we prove

THEOREM 12. *$B^* \in [\sqrt{2}, 2)$. $\zeta(B) > 0$ on $[0, B^*)$, and $\zeta(B) < 0$ on $(B^*, 2]$.*

We also complete the proof of Proposition 1. We begin by proving some properties of the derivatives of $\frac{D_x(y)}{D_{x-1}(y)}$, which will then enable us to analyze $\frac{d}{dB}\zeta(B)$ using the multivariate chain rule.

LEMMA 18.

(i) $D_{x-1}(y) \in (0, \infty)$ for $x \leq 1$ and all y.

(ii) $\frac{D_x(y)}{D_{x-1}(y)}$ is a smooth function of y for $x \leq 1$, and

$$\frac{d}{dy}\frac{D_x(y)}{D_{x-1}(y)} = x + \Big(\frac{D_x(y)}{D_{x-1}(y)}\Big)^2 - y\frac{D_x(y)}{D_{x-1}(y)}. \tag{63}$$

(iii) $\frac{D_x(y)}{D_{x-1}(y)}$ is a smooth function of x for $x \leq 1$ and all y.

PROOF. Note that if $x \leq 1$ then $D_{x-1}(y) > 0$ for all y by (4), from which assertion i. follows. We now prove assertion ii. Since $D_x(y)$ is an entire function of y for each fixed x, it follows from i. that for $x \leq 1$, $\frac{D_x(y)}{D_{x-1}(y)}$ is a smooth function of y, and

$$\frac{d}{dy}\frac{D_x(y)}{D_{x-1}(y)}$$

$$
\begin{aligned}
&= \frac{D_{x-1}(y)\frac{d}{dy}D_x(y) - D_x(y)\frac{d}{dy}D_{x-1}(y)}{D^2_{x-1}(y)} \\
&= \frac{D_{x-1}(y)\big(xD_{x-1}(y) - \frac{1}{2}yD_x(y)\big) - D_x(y)\big((x-1)D_{x-2}(y) - \frac{1}{2}yD_{x-1}(y)\big)}{D^2_{x-1}(y)} \quad \text{by (23)} \\
&= \frac{xD^2_{x-1}(y) - \frac{1}{2}yD_x(y)D_{x-1}(y) - (x-1)D_x(y)D_{x-2}(y) + \frac{1}{2}yD_x(y)D_{x-1}(y)}{D^2_{x-1}(y)} \\
&= x - \frac{(x-1)D_x(y)D_{x-2}(y)}{D^2_{x-1}(y)} \\
&= x - \frac{(x-1)D_x(y)(\frac{D_x(y)-yD_{x-1}(y)}{1-x})}{D^2_{x-1}(y)} \quad \text{by (22)} \\
&= x + \frac{D_x(y)(D_x(y) - yD_{x-1}(y))}{D^2_{x-1}(y)} \\
&= x + (\frac{D_x(y)}{D_{x-1}(y)})^2 - y\frac{D_x(y)}{D_{x-1}(y)}.
\end{aligned}
$$

Also, assertion iii. follows from the fact that $D_x(y)$ is an entire function of x for all y, and (4). $\square$

We now prove some properties of $\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$ by applying the multivariate chain rule to Lemma 18.

Lemma 19.

(i) $D_{\frac{B^2}{4}-1}(-B) \in (0,\infty)$ *on* $(-\infty, 2]$.

(ii) $\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$ *is a differentiable function of* B *on* $(-\infty, 2]$, *and on* $(-\infty, 2]$ *we have*

$$
\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} = \frac{B}{2}\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{\substack{x=\frac{B^2}{4}\\ y=-B}} - \frac{B^2}{4} - (\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)})^2 - B\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}.
$$

Proof. $D_{\frac{B^2}{4}-1}(-B) > 0$ for $B \le 2$ by (4). Thus we may apply the multivariate chain rule to find that $\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$ is a differentiable function of B on $(-\infty, 2]$, and

$$
\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}
$$

$$
\begin{aligned}
&= \frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{\substack{x=\frac{B^2}{4}\\ y=-B}}\frac{d}{dB}\frac{B^2}{4} + \frac{d}{dy}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{\substack{x=\frac{B^2}{4}\\ y=-B}}\frac{d}{dB}(-B)\\
&= \frac{B}{2}\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{\substack{x=\frac{B^2}{4}\\ y=-B}} - \frac{d}{dy}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{\substack{x=\frac{B^2}{4}\\ y=-B}}\\
&= \frac{B}{2}\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{\substack{x=\frac{B^2}{4}\\ y=-B}} - \frac{B^2}{4} - \left(\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}\right)^2 - B\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} \quad \text{by Lemma 18.ii.}
\end{aligned}
$$

□

Although it seems clear from Figure 1 that $\zeta(B)$ is strictly negative on $(B^*, 2)$, the formal proof of this fact is somewhat involved, and will necessitate careful arguing about $\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$. Due to the fact that $\frac{d}{dx}D_x(y)$ has no simple analytic form $\big($ as opposed to $\frac{d}{dy}D_x(y)$, see (23) $\big)$, to proceed we will have to derive good bounds for $\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}$. Our bounds will rely on the concavity of certain functions. We begin by proving that

Lemma 20. *For any fixed $y < 0$, $\frac{D_x(y)}{D_{x-1}(y)}$ is a concave function of x on $(0,1)$.*

Proof. For any fixed $B > 0$, $z_\infty(x)$ is a concave function of x on $(0,1)$, which follows from Lemma 2, Proposition 2, and the fact that pointwise limits of concave functions are concave. The lemma follows, since $\frac{D_x(-B)}{D_{x-1}(-B)} = z_\infty(x) - B$ (by Proposition 2). □

We now use the concavity of $\frac{D_x(y)}{D_{x-1}(y)}$ and the associated monotonicity of $\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}$ to prove that

Lemma 21. *For any $y < 0$ and $x_0 \in (0,1)$,*

$$\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{x=x_0} < \frac{1}{x_0}\frac{D_{x_0}(y)}{D_{x_0-1}(y)}.$$

Proof. By Lemma 18.iii and the Mean Value Theorem, there exists $c_{x_0,y} \in (0, x_0)$ such that

$$\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{x=c_{x_0,y}} = \frac{\frac{D_{x_0}(y)}{D_{x_0-1}(y)} - \frac{D_0(y)}{D_{-1}(y)}}{x_0}.$$

Thus

$$
\begin{aligned}
(64)\qquad \frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{x=x_0} &\le \frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}\bigg|_{x=c_{x_0,y}} \quad \text{by Lemma 20 and the def. of concavity}\\
&= \frac{\frac{D_{x_0}(y)}{D_{x_0-1}(y)} - \frac{D_0(y)}{D_{-1}(y)}}{x_0}\\
&< \frac{1}{x_0}\frac{D_{x_0}(y)}{D_{x_0-1}(y)} \quad \text{since } \frac{D_0(y)}{D_{-1}(y)} > 0 \text{ by (3), and } x_0 > 0.
\end{aligned}
$$

□

We now plug our bounds on $\frac{d}{dx}\frac{D_x(y)}{D_{x-1}(y)}$ from Lemma 21 into our bounds on $\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$ from Lemma 19 to prove that

Lemma 22. *For $0 < B < 2$,*

$$\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} < (\frac{2}{B} - B)\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} - \frac{B^2}{4} - (\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)})^2 \leq \frac{1}{B^2} - 1.$$

Proof. By Lemma 19.ii and Lemma 21, for $0 < B < 2$,

$$\begin{aligned}\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} &< \frac{B}{2}\frac{1}{\frac{B^2}{4}}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} - \frac{B^2}{4} - (\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)})^2 - B\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} \\ &= (\frac{2}{B} - B)\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} - \frac{B^2}{4} - (\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)})^2.\end{aligned}$$

This proves the first inequality. Furthermore, it follows that there exists $x_B \in \mathbb{R}$ (we may take $x_B = \frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$) such that $\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} \leq (\frac{2}{B} - B)x_B - x_B^2 - \frac{B^2}{4}$. Since by elementary calculus the function (of x) $(\frac{2}{B} - B)x - x^2 - \frac{B^2}{4}$ attains its maximum at $x = \frac{(\frac{2}{B}-B)}{2}$, we find that for all $B \in (0, 2)$,

$$\begin{aligned}\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} &\leq (\frac{2}{B} - B)(\frac{(\frac{2}{B} - B)}{2}) - (\frac{(\frac{2}{B} - B)}{2})^2 - \frac{B^2}{4} \\ &= \frac{1}{B^2} - 1.\end{aligned}$$

□

We are now in a position to study the zeros of $\zeta(B)$.

Lemma 23. *$\zeta(B) > 0$ on $[0, B^*)$, and $B^* \in (0, 2)$.*

Proof. By (3),

$$\zeta(0) = \frac{D_0(0)}{D_{-1}(0)} = \frac{1}{\sqrt{2}\int_0^\infty e^{-t^2}dt} > 0, \tag{65}$$

and

$$\zeta(2) = \frac{D_1(-2)}{D_0(-2)} + 1 = \frac{-2e^{-1}}{e^{-1}} + 1 = -1 < 0. \tag{66}$$

By Lemma 19.ii and definitions, $\zeta(B)$ is a differentiable (and continuous) function of B on $(-\infty, 2]$. It follows from the Intermediate Value Theorem that $B^* \in (0, 2)$, and $\zeta(B) > 0$ on $[0, B^*)$. □

Lemma 24. *$B^* \in [\sqrt{2}, 2)$.*

Proof. It is proven in [40] (Theorem 4.1, i.) that

$$\gamma_n \geq \inf_{k\geq 1} \Big(\lambda_n + \min(k,n) - \sqrt{\lambda_n}\big(\sqrt{\min(k-1,n)} + \sqrt{\min(k,n)}\big)\Big). \tag{67}$$

Note that for $1 \leq k \leq n$,

$$\begin{aligned} \lambda_n + \min(k,n) - \sqrt{\lambda_n}\big(\sqrt{\min(k-1,n)} + \sqrt{\min(k,n)}\big) &= \lambda_n + k - \sqrt{\lambda_n}(\sqrt{k-1} + \sqrt{k}) \\ &= (\sqrt{\lambda_n} - \sqrt{k})^2 + \sqrt{\lambda_n}(\sqrt{k} - \sqrt{k-1}) \\ &= (\sqrt{\lambda_n} - \sqrt{k})^2 + \frac{\sqrt{\lambda_n}}{\sqrt{k} + \sqrt{k-1}} \\ &\geq \frac{1}{2}\frac{\sqrt{\lambda_n}}{\sqrt{k}} \\ &\geq \frac{1}{2}\frac{\sqrt{\lambda_n}}{\sqrt{n}}. \end{aligned} \tag{68}$$

For all $k \geq n+1$, the right-hand side of (67) equals $(\sqrt{n} - \sqrt{\lambda_n})^2$. Combining this fact with (67) and (68), we find that

$$\gamma_n \geq \min\big(\frac{1}{2}\frac{\sqrt{\lambda_n}}{\sqrt{n}}, (\sqrt{n} - \sqrt{\lambda_n})^2\big). \tag{69}$$

By Lemma 4, $\lim_{n\to\infty}(\sqrt{n} - \sqrt{\lambda_n})^2 = \frac{B^2}{4}$. Trivially, $\lim_{n\to\infty} \frac{1}{2}\frac{\sqrt{\lambda_n}}{\sqrt{n}} = \frac{1}{2}$. Thus for any $B < \sqrt{2}$ and all sufficiently large n, $\frac{1}{2}\frac{\sqrt{\lambda_n}}{\sqrt{n}} > (\sqrt{n} - \sqrt{\lambda_n})^2$. It follows that $\gamma_n \geq (\sqrt{n} - \sqrt{\lambda_n})^2$ for all sufficiently large n by (69).

Now, suppose for contradiction that $B^* < \sqrt{2}$. By Lemma 22 and the fact that $B^* \in (0,2)$ by Lemma 23, we have that $\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}$ is a differentiable function of B at B^*, and

$$\frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)}\bigg|_{B=B^*}$$

$$\begin{aligned} &< (\frac{2}{B^*} - B^*)(-\frac{B^*}{2}) - \frac{(B^*)^2}{4} - (-\frac{B^*}{2})^2 \quad \text{since by def.}\ \frac{D_{\frac{(B^*)^2}{4}}(-B^*)}{D_{\frac{(B^*)^2}{4}-1}(-B^*)} = -\frac{B^*}{2} \\ &= -1. \end{aligned}$$

It follows from definitions and the fact that $\frac{d}{dB}\frac{B}{2} = \frac{1}{2}$ that

$$\frac{d}{dB}\zeta(B)\bigg|_{B=B^*} < 0.$$

Thus there exists $\epsilon > 0$ s.t. $\zeta(B) < 0$ on $(B^*, B^* + \epsilon)$ since $\zeta(B^*) = 0$. It follows that there exists $B' \in (0, \sqrt{2})$ such that $\zeta(B') < 0$. Thus if we define all relevant functions in terms of B', $\Psi_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) < 0$ for all sufficiently large n by Corollary 2. Also, $(\sqrt{n} - \sqrt{\lambda_n})^2 < 1$ for all sufficiently large n by Lemma 4. Thus by Lemma 3.i, for all sufficiently large n, $\gamma_n < (\sqrt{n} - \sqrt{\lambda_n})^2$. But this is a contradiction, since we have already shown that $B' < \sqrt{2}$ implies that $\gamma_n \geq (\sqrt{n} - \sqrt{\lambda_n})^2$ for all sufficiently large n. □

Proof of Theorem 12. In light of Lemma 23 and Lemma 24, all that remains to be shown is that $\zeta(B) < 0$ on $(B^*, 2)$. In light of Lemma 24, it would suffice to prove that $\zeta(B)$ is strictly decreasing on $(\sqrt{2}, 2)$. But for $B \in (\sqrt{2}, 2)$,

$$\begin{aligned}\frac{d}{dB}\zeta(B) &= \frac{d}{dB}\frac{D_{\frac{B^2}{4}}(-B)}{D_{\frac{B^2}{4}-1}(-B)} + \frac{d}{dB}\frac{B}{2} \\ &< \frac{1}{B^2} - 1 + \frac{1}{2} \quad \text{by Lemma 22} \\ &= 0.\end{aligned}$$

This demonstrates that $\zeta(B)$ is strictly decreasing on $(\sqrt{2}, 2)$, concluding the proof of Theorem 12. $\square$

Proof of Proposition 1. That $B^* \in [\sqrt{2}, 2)$ follows immediately from Theorem 12. That $Z^+(\Psi_\infty) < \min(1, \frac{B^2}{4})$ for $B > B^*$ follows immediately from Theorem 12 and Lemma 17.i. $\square$

7. Proof of Theorem 5, Corollary 1, and Theorem 6.

Proof of Theorem 5. First, suppose $0 < B < B^*$. Then $B < 2$ by Theorem 12, and thus $(\sqrt{n} - \sqrt{\lambda_n})^2 < 1$ for all sufficiently large n by Lemma 4. $\zeta(B) > 0$ by Theorem 12, and thus since $B < 2$, $\Psi_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) > 0$ for all sufficiently large n by Corollary 2. It follows from Lemma 3.ii that $\gamma_n = (\sqrt{n} - \sqrt{\lambda_n})^2$ for all sufficiently large n. That $\lim_{n\to\infty} \gamma_n = \frac{B^2}{4}$ then follows from Lemma 4.

Now, suppose $B = B^*$. Since $B^* < 2$ by Theorem 12, it follows from Lemma 4 that $(\sqrt{n} - \sqrt{\lambda_n})^2 < 1$ for all sufficiently large n. By Lemma 3(i and ii), for all sufficiently large n, either $\Psi_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) < 0$, in which case $\gamma_n = Z^+(\Psi_n)$ is the unique zero of Ψ_n in the interval $\big(0, (\sqrt{n} - \sqrt{\lambda_n})^2\big)$, or $\gamma_n = (\sqrt{n} - \sqrt{\lambda_n})^2$. Let $\{n_i\}$ denote the subsequence of $\{n\}$ for which $\Psi_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) < 0$. If $\{n_i\}$ is a finite set, then trivially $\gamma_n = (\sqrt{n} - \sqrt{\lambda_n})^2$ for all sufficiently large n. That $\lim_{n\to\infty} \gamma_n = \frac{(B^*)^2}{4}$ then follows from Lemma 4.

Alternatively, suppose $\{n_i\}$ is an infinite set. Suppose for contradiction that $\liminf_{i\to\infty} Z^+(\Psi_{n_i}) < \frac{(B^*)^2}{4}$. Note that by Lemma 17.ii, $Z^+(\Psi_\infty) = \frac{(B^*)^2}{4}$, and $\Psi_\infty(x) > 0$ on $[0, \frac{(B^*)^2}{4})$. It follows that there exists $\epsilon > 0$ such that $0 < \liminf_{i\to\infty} Z^+(\Psi_{n_i}) + \epsilon < \frac{(B^*)^2}{4} < 1$ and $\Psi_\infty(\liminf_{i\to\infty} Z^+(\Psi_{n_i}) + \epsilon) > 0$. It follows from Lemma 4 and Theorem 10 that for all sufficiently large i, $0 < \liminf_{i\to\infty} Z^+(\Psi_{n_i}) + \epsilon < (\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2 < 1$ and $\Psi_{n_i}(\liminf_{i\to\infty} Z^+(\Psi_{n_i}) + \epsilon) > 0$. By the monotonicity of $\Psi_{n_i}(x)$ (from Lemma 1.iv), it follows that for all sufficiently large i, $\Psi_{n_i}(x) > 0$ on $(-\infty, \liminf_{i\to\infty} Z^+(\Psi_{n_i}) + \epsilon)$. But by the definition of $\liminf$, there exists an infinite strictly increasing sequence of integers $\{n_i'\}$ s.t. $Z^+(\Psi_{n_i'}) < \liminf_{i\to\infty} Z^+(\Psi_{n_i}) + \epsilon$ for all i. Thus for all sufficiently large i, $\Psi_{n_i'}(Z^+(\Psi_{n_i'})) > 0$. This is a contradiction, since $\Psi_{n_i'}(Z^+(\Psi_{n_i'})) = 0$ for all i. This proves that $\liminf_{i\to\infty} Z^+(\Psi_{n_i}) \geq \frac{(B^*)^2}{4}$.

Note that $(\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2 < 1$ for all sufficiently large i by Lemma 4 and Proposition 1. It follows from Lemma 3.i that for all sufficiently large i, $Z^+(\Psi_{n_i}) < (\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2$, and thus $\limsup_{i\to\infty} Z^+(\Psi_{n_i}) \leq \frac{(B^*)^2}{4}$ by Lemma 4. Combining the above, we find that $\lim_{i\to\infty} Z^+(\Psi_{n_i}) = \frac{(B^*)^2}{4}$. Thus γ_n alternates between two sequences (that of the $(\sqrt{n} - \sqrt{\lambda_n})^2$ and that of the $Z^+(\Psi_{n_i})$), both of which converge to $\frac{(B^*)^2}{4}$, proving that $\lim_{n\to\infty} \gamma_n = \frac{(B^*)^2}{4} = Z^+(\Psi_\infty)$.

Now, suppose $B \in (B^*, 2)$. Then $(\sqrt{n} - \sqrt{\lambda_n})^2 < 1$ for all sufficiently large n by Lemma 4.

$\zeta(B) < 0$ by Theorem 12, and thus since $B < 2$, $\Psi_n\big((\sqrt{n} - \sqrt{\lambda_n})^2\big) < 0$ for all sufficiently large n by Corollary 2. It follows from Lemma 3.i that $\gamma_n = Z^+(\Psi_n)$ for all sufficiently large n. That $\lim_{n\to\infty} \gamma_n = Z^+(\Psi_\infty)$ then follows from Theorem 11.

Now, suppose $B \geq 2$. Then $(\sqrt{n} - \sqrt{\lambda_n})^2 > 1$ for all sufficiently large n by Lemma 4. It follows from Lemma 3.iii that $\gamma_n = Z^+(\Psi_n)$ for all sufficiently large n. That $\lim_{n\to\infty} \gamma_n = Z^+(\Psi_\infty)$ then follows from Theorem 11. This treats all cases, and we conclude that $\lim_{n\to\infty} \gamma_n \overset{\Delta}{=} \gamma_B$ exists for all $B > 0$. We also conclude that for $0 < B \leq B^*$, $\gamma_B = \frac{B^2}{4}$; for $B \geq B^*$, $\gamma_B = Z^+(\Psi_\infty)$. ☐

PROOF OF COROLLARY 1. Suppose for contradiction that $\liminf_{n\to\infty} \sqrt{n}(1 - \rho_n^*) < B^*$. Then there exists $\epsilon > 0$ and an infinite strictly increasing sequence of integers $\{n_i\}$ s.t. $\sqrt{n_i}(1 - \rho_{n_i}^*) < B^* - \epsilon$, and thus $\rho_{n_i}^* > 1 - \frac{B^* - \epsilon}{\sqrt{n_i}}$. Consider the sequence $\{Z_i\}$ of continuous time markov chains, in which Z_i is an $M/M/n_i$ queueing system with $\lambda_{n_i} = n_i - (B^* - \epsilon)\sqrt{n_i}$, $\mu = 1$. Since $B^* - \epsilon < B^*$, it follows from Theorem 12 that $\zeta(B^* - \epsilon) > 0$ and $B^* - \epsilon < 2$. Now, let us define all relevant functions (i.e. $\Psi_{n_i}(x)$) w.r.t. $B^* - \epsilon$. It then follows from Corollary 2 and Lemma 4 that for all sufficiently large i, $\Psi_{n_i}\big((\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2\big) > 0$, and $(\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2 < 1$. Thus by Lemma 3.ii, $\gamma_{n_i} = (\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2$ for all sufficiently large i. However, note that by assumption $\frac{\lambda_{n_i}}{n_i\mu} = 1 - \frac{B^* - \epsilon}{\sqrt{n_i}} < \rho_{n_i}^*$ for all i. But this is a contradiction, since by Theorem 4, $\frac{\lambda_{n_i}}{n_i\mu} < \rho_{n_i}^*$ implies that the spectral gap γ_{n_i} of Z_i is strictly less than $(\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2$.

Alternatively, suppose for contradiction that $\limsup_{n\to\infty} \sqrt{n}(1 - \rho_n^*) > B^*$. Then there exists $\epsilon \in (0, 2 - B^*)$ and an infinite strictly increasing sequence of integers $\{n_i\}$ s.t. $\sqrt{n_i}(1 - \rho_{n_i}^*) > B^* + \epsilon$, and thus $\rho_{n_i}^* < 1 - \frac{B^* + \epsilon}{\sqrt{n_i}}$. Consider the sequence $\{Z_i\}$ of continuous time Markov chains, in which Z_i is an $M/M/n_i$ queueing system with $\lambda_{n_i} = n_i - (B^* + \epsilon)\sqrt{n_i}$, $\mu = 1$. Since $B^* + \epsilon \in (B^*, 2)$, it follows from Theorem 12 that $\zeta(B^* + \epsilon) < 0$. Now, let us define all relevant functions w.r.t. $B^* + \epsilon$. It then follows from Corollary 2 and Lemma 4 that for all sufficiently large i, $\Psi_{n_i}\big((\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2\big) < 0$, and $(\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2 < 1$. Thus by Lemma 3.i, $\gamma_{n_i} = Z^+(\Psi_{n_i}) < (\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2$ for all sufficiently large i. However, note that by assumption $\frac{\lambda_{n_i}}{n_i\mu} = 1 - \frac{B^* + \epsilon}{\sqrt{n_i}} > \rho_{n_i}^*$ for all i. But this is a contradiction, since by Theorem 4, $\frac{\lambda_{n_i}}{n_i\mu} > \rho_{n_i}^*$ implies that the spectral gap γ_{n_i} of Z_i is equal to $(\sqrt{n_i} - \sqrt{\lambda_{n_i}})^2$. It follows that $\liminf_{n\to\infty} \sqrt{n}(1 - \rho_n^*) = \limsup_{n\to\infty} \sqrt{n}(1 - \rho_n^*) = B^*$. ☐

We now prove a series of lemmas, which will collectively prove Theorem 6. To make the dependence on B clear, let $\Psi_\infty(B, x)$ denote $\Psi_\infty(x)$ defined w.r.t. B.

LEMMA 25. *For each $B > 0$ there exists $\epsilon_B > 0$ s.t. $0 < B' - B < \epsilon_B$ implies $\gamma_{B'} > \gamma_B$.*

PROOF. Suppose $0 < B < B^*$. Then $\gamma_B = \frac{B^2}{4}$ by Theorem 5, and thus trivially γ_B is strictly increasing on $(0, B^*)$.

Suppose $B \geq B^*$. By Proposition 1 and Theorem 5,

$$\text{For all } B > B^*, 0 < \gamma_B < \min(1, \frac{B^2}{4}). \text{ Also, } \; 0 < \gamma_{B^*} = \frac{(B^*)^2}{4} < 1. \tag{70}$$

It follows from (70), Lemma 18.ii, Theorem 5, and the definitions of $\Psi_\infty(x)$ and γ_B that for all

$B \geq B^*$,

$$\begin{aligned}
\frac{d}{dy}\frac{D_x(-y)}{D_{x-1}(-y)}\bigg|_{\substack{x=\gamma_B\\ y=B}} &= -\gamma_B - (\frac{D_{\gamma_B}(-B)}{D_{\gamma_B-1}(-B)})^2 - B\frac{D_{\gamma_B}(-B)}{D_{\gamma_B-1}(-B)} \\
&= -\gamma_B - \Big(-\frac{1}{2}\big(B+(B^2-4\gamma_B)^{\frac{1}{2}}\big)\Big)^2 - B\Big(-\frac{1}{2}\big(B+(B^2-4\gamma_B)^{\frac{1}{2}}\big)\Big) \\
&= -\gamma_B + \Big(\frac{1}{2}\big(B+(B^2-4\gamma_B)^{\frac{1}{2}}\big)\Big)\Big(\frac{1}{2}\big(B-(B^2-4\gamma_B)^{\frac{1}{2}}\big)\Big) \\
&= -\gamma_B + \frac{4\gamma_B}{4} \\
&= 0. \qquad (71)
\end{aligned}$$

It follows from (70) that for all $B \geq B^*$ and $\epsilon > 0$,

$$\frac{1}{2}\Big((B+\epsilon)+\big((B+\epsilon)^2-4\gamma_B\big)^{\frac{1}{2}}\Big) - \frac{1}{2}\Big(B+(B^2-4\gamma_B)^{\frac{1}{2}}\Big) > \frac{\epsilon}{2}. \qquad (72)$$

Thus in light of the definition of $\Psi_\infty(x)$, it follows from (71), (72), and the definition of derivative that for $B \geq B^*$, there exists $\epsilon_B > 0$ such that $B' \in (B, B+\epsilon_B)$ implies $\Psi_\infty(B', \gamma_B) > 0$. The lemma then follows from Lemma 17.i and Theorem 5. ☐

Lemma 26. *γ_B is continuous on $(0, \infty)$.*

Proof. First, suppose $B \in (0, B^*]$. Then $\gamma_B = \frac{B^2}{4}$ by Theorem 5, from which it follows trivially that γ_B is continuous on $(0, B^*)$ and left-continuous at B^*.

Now, suppose $B > B^*$. It follows from (70) that there exists $\delta_B \in \big(0, \min(1-\gamma_B, \frac{B^2}{4}-\gamma_B, \gamma_B)\big)$ and $\delta_{0,B} \in \big(0, \min(B, B-B^*)\big)$ s.t. $\gamma_B + \delta_B < \min\big(1, \frac{(B-\delta_{0,B})^2}{4}\big)$ and $B - \delta_{0,B} > B^*$. It follows from Lemma 16 that $|B'-B| < \delta_{0,B}$ implies $\Psi_\infty(B', x)$ is a differentiable function of x on $\big(\gamma_B - \delta_B, \gamma_B + \delta_B\big)$, and for $x \in \big(\gamma_B - \delta_B, \gamma_B + \delta_B\big)$,

$$\begin{aligned}
\frac{d}{dx}\Psi_\infty(B', x) &\leq -\big((B')^2-4x\big)^{-1} \\
&\leq -\big((B')^2\big)^{-1} \text{ since } \ x > 0 \\
&\leq -\big(B+\delta_{0,B}\big)^{-2} \\
&< -(B+B)^{-2} \text{ since } \delta_{0,B} < B \\
&= -(4B^2)^{-1}. \qquad (73)
\end{aligned}$$

By (70) and the continuity of $\frac{D_x(-y)}{D_{x-1}(-y)}$ (as a function of y) ensured by Lemma 18.ii, for all $\epsilon > 0$ there exists $\delta_{1,B,\epsilon} > 0$ s.t. $|B'-B| < \delta_{1,B,\epsilon}$ implies

$$\left|\frac{D_{\gamma_B}(-B)}{D_{\gamma_B-1}(-B)} - \frac{D_{\gamma_B}(-B')}{D_{\gamma_B-1}(-B')}\right| < \frac{\epsilon}{8B^2}. \qquad (74)$$

Note that for $x > 0$, $\frac{1}{2}\big(B+(B^2-4x)^{\frac{1}{2}}\big)$ is a continuous function of B on $(2x^{\frac{1}{2}}, \infty)$, and a strictly increasing function of B on $[2x^{\frac{1}{2}}, \infty)$. It follows from (70) that for all $\epsilon > 0$ there exists $\delta_{2,B,\epsilon} > 0$

s.t. $|B' - B| < \delta_{2,B,\epsilon}$ implies $B' > 2\gamma_B^{\frac{1}{2}}$, and

$$\left| \frac{1}{2}\big(B + (B^2 - 4\gamma_B)^{\frac{1}{2}}\big) - \frac{1}{2}\big(B' + ((B')^2 - 4\gamma_B)^{\frac{1}{2}}\big) \right| < \frac{\epsilon}{8B^2}. \tag{75}$$

From the definition of $\Psi_\infty(x)$, (74),(75), the triangle inequality, and the fact that $\Psi_\infty(B, \gamma_B) = 0$, for all $\epsilon > 0$ there exists $\delta_{3,B,\epsilon} > 0$ s.t. $|B' - B| < \delta_{3,B,\epsilon}$ implies

$$|\Psi_\infty(B', \gamma_B)| < \frac{\epsilon}{4B^2}. \tag{76}$$

It follows from (73), (76), and the construction of δ_B that for all $\epsilon \in (0, \delta_B)$, $|B'-B| < \min(\delta_{0,B}, \delta_{3,B,\epsilon})$ implies $\Psi_\infty(B', x)$ has a zero in $(\gamma_B - \epsilon, \gamma_B + \epsilon)$. Thus since $(\gamma_B - \epsilon, \gamma_B + \epsilon)$ is itself contained in the interval $\big(0, \min(1, \frac{B^2}{4})\big)$ by construction of δ_B, it follows from Lemma 17.i and Theorem 12 that this zero must be $Z^+(\Psi_\infty)$ (defined w.r.t. B'). Thus by Theorem 5, $|B' - B| < \min(\delta_{0,B}, \delta_{3,B,\epsilon})$ implies $|\gamma_B - \gamma_{B'}| < \epsilon$. This concludes the proof that γ_B is continuous on (B^*, ∞).

We now prove that γ_B is right-continuous at B^*. By Lemma 25, there exists $\delta > 0$ s.t. $B' \in (B^*, B^*+\delta)$ implies $\gamma_{B'} > \gamma_{B^*}$. It follows from (70) that for all $B' \in (B^*, B^*+\delta)$, $\frac{(B^*)^2}{4} < \gamma_B < \frac{(B')^2}{4}$; the right-continuity of γ_B at B^* then follows from the continuity of $\frac{B^2}{4}$. Combining the above, we find that γ_B is continuous on $(0, \infty)$. ☐

Corollary 3. *γ_B is strictly increasing on* $(0, \infty)$.

Proof. Suppose $0 < B_1 < B_2$. Let $\gamma_{\max} = \sup_{B \in [B_1, B_2]} \gamma_B$. Since γ_B is continuous and non-negative on $[B_1, B_2]$ we know that $0 < \gamma_B < \infty$, and $\gamma_{\max}$ is attained by at least one $B \in [B_1, B_2]$. Now, suppose for contradiction that there exists some $B_{\max} \in [B_1, B_2)$ attaining this maximum. Then by Lemma 25, there exists $\epsilon_{B_{\max}} > 0$ s.t. $B' \in (B_{\max}, B_{\max} + \epsilon_{B_{\max}})$ implies $B' \in [B_1, B_2]$ and $\gamma_{B'} > \gamma_{B_{\max}} = \gamma_{\max}$. But this is a contradiction, since by construction $\gamma_{\max} = \sup_{B \in [B_1, B_2]} \gamma_B$. It follows that the unique $B \in [B_1, B_2]$ at which γ_B is maximized is B_2, and thus $\gamma_{B_2} > \gamma_{B_1}$. ☐

Lemma 27. $B \geq 10$ *implies* $0 < 1 - e^{-\frac{(B-1)^2}{2}}, 1 - e^{-\frac{B^2}{2} - 8\log(3B+7)} < \min(1, \frac{B^2}{4})$, $\Psi_\infty(1 - e^{-\frac{(B-1)^2}{2}}) > 0$, *and* $\Psi_\infty(1 - e^{-\frac{B^2}{2} - 8\log(3B+7)}) < 0$.

Proof. We proceed by bounding the terms that appear in the representation for $\Psi_\infty(x)$ when one expands the parabolic cylinder functions using their integral representations, as in Lemma 15. Suppose $B \geq 10$, and $.5 < x < 1$. Then

$$\int_0^\infty e^{-\frac{(B-z)^2}{2}} z^{1-x} dz$$

$$\begin{aligned}
&\leq \int_0^{B+1} z^{1-x} dz + \int_{B+1}^\infty e^{-\frac{(z-B)^2}{2}} z dz \quad \text{since } e^{-\frac{(B-z)^2}{2}} < 1 \text{ , and } z^{1-x} < z \text{ for } z > B+1 \\
&\leq \int_0^{B+1} z^{1-x} dz + \int_{B+1}^\infty e^{-\frac{(z-B)}{2}} z dz \quad \text{since } 0 < z - B < (z-B)^2 \text{ for } z > B+1 \\
&= \frac{(B+1)^{2-x}}{2-x} + 2e^{-\frac{1}{2}}(3+B) \\
&\qquad \text{since } \int_0^{B+1} z^{1-x} dz = \frac{(B+1)^{2-x}}{2-x}, \text{ and } \int_{B+1}^\infty e^{-\frac{(z-B)}{2}} z dz = 2e^{-\frac{1}{2}}(3+B)
\end{aligned}$$

$$\leq (B+1)^2 + 6 + 2B \quad \leq \quad (3B+7)^2 \quad \text{since } 0 < x < 1. \tag{77}$$

Similarly,

$$\begin{aligned}\int_0^\infty e^{-\frac{(B-z)^2}{2}} z^{1-x}dz &\geq \int_B^\infty e^{-\frac{(B-z)^2}{2}}dz \quad \text{by non-negativity, and since } z^{1-x} > 1 \text{ for } \; z > B \\ &= \sqrt{\frac{\pi}{2}} \; > 1 \quad \text{by standard integration techniques.} \end{aligned} \tag{78}$$

Likewise,

$$\begin{aligned}\int_0^\infty e^{-\frac{(B-z)^2}{2}} z^{-x}dz &= \int_0^1 e^{-\frac{(B-z)^2}{2}} z^{-x}dz + \int_1^\infty e^{-\frac{(B-z)^2}{2}} z^{-x}dz \\ &\leq e^{-\frac{(B-1)^2}{2}} \int_0^1 z^{-x}dz + \int_1^\infty e^{-\frac{(B-z)^2}{2}} z^{-x}dz \;\; \text{since } B \geq 10 \text{ implies } e^{-\frac{(B-z)^2}{2}} \text{ inc. on } [0,1] \\ &\leq e^{-\frac{(B-1)^2}{2}} \int_0^1 z^{-x}dz + \int_1^\infty e^{-\frac{(B-z)^2}{2}} dz \quad \text{since } \; z^{-x} \leq 1 \text{ for } \; 0 < x < 1, z \geq 1 \\ &\leq e^{-\frac{(B-1)^2}{2}} \int_0^1 z^{-x}dz + \int_{-\infty}^\infty e^{-\frac{(B-z)^2}{2}} dz \quad \text{by non-negativity} \\ &= \frac{e^{-\frac{(B-1)^2}{2}}}{1-x} + \sqrt{2\pi} \;\; \leq \;\; \frac{e^{-\frac{(B-1)^2}{2}}}{1-x} + 3 \quad \text{by standard integration techniques.} \end{aligned} \tag{79}$$

Also,

$$\begin{aligned}\int_0^\infty e^{-\frac{(B-z)^2}{2}} z^{-x}dz &\geq \int_0^1 e^{-\frac{(B-z)^2}{2}} z^{-x}dz \\ &\geq e^{-\frac{B^2}{2}} \int_0^1 z^{-x}dz \quad \text{since } B \geq 10 \text{ implies } e^{-\frac{(B-z)^2}{2}} \text{ increasing on } [0,1] \\ &= \frac{e^{-\frac{B^2}{2}}}{1-x}. \end{aligned} \tag{80}$$

It follows from (77) - (80) and Lemma 15 that

$$\Big(\frac{e^{-\frac{(B-1)^2}{2}}}{1-x} + 3\Big)^{-1} \leq \frac{D_x(-B)}{D_{x-1}(-B)} + B \leq (3B+7)^2 e^{\frac{B^2}{2}}(1-x). \tag{81}$$

Note that $.5 < x < 1$ and $B \geq 10$ implies $x < \frac{B^2}{4}$, and

$$\frac{1}{2}\big((B^2-4x)^{\frac{1}{2}} - B\big) = -2x\big((B^2-4x)^{\frac{1}{2}} + B\big)^{-1}. \tag{82}$$

It follows that the left-hand side of (82) is at least $-2B^{-1}$, since $x < 1$ and $(B^2-4x)^{\frac{1}{2}} + B > B$. Also, the left-hand side of (82) is at most $-.5B^{-1}$, since $x > .5$ and $(B^2-4x)^{\frac{1}{2}} + B < 2B$ for $B \geq 10$ and $x < 1$. Combining with (81) and the definition of $\Psi_\infty(x)$, we find that

$$\Big(\frac{e^{-\frac{(B-1)^2}{2}}}{1-x} + 3\Big)^{-1} - 2B^{-1} \leq \Psi_\infty(x) \leq (3B+7)^2 e^{\frac{B^2}{2}}(1-x) - .5B^{-1}. \tag{83}$$

Since $B \geq 10$ implies $1 - e^{-\frac{B^2}{2} - 8\log(3B+7)} \in (.5, 1)$, it follows from (83) that

$$\begin{aligned}
\Psi_\infty(1 - e^{-\frac{B^2}{2} - 8\log(3B+7)}) &\leq (3B+7)^2 e^{\frac{B^2}{2}} e^{-\frac{B^2}{2} - 8\log(3B+7)} - .5B^{-1} \\
&= (3B+7)^{-6} - .5B^{-1} \\
&< 0 \quad \text{since } (3B+7)^{-6} < B^{-6}\ ,\ B^{-6} - .5B^{-1} < 0 \text{ for } B \geq 10. \qquad (84)
\end{aligned}$$

Similarly,

$$\Psi_\infty(1 - e^{-\frac{(B-1)^2}{2}}) \geq \frac{1}{4} - 2B^{-1} > 0 \text{ for } B > 10. \qquad (85)$$

The lemma follows by combining (84) and (85). ☐

PROOF OF THEOREM 6. That γ_B is a continuous, strictly increasing function of B follows immediately from Lemma 26 and Corollary 3. That $\lim_{B\to\infty} \gamma_B = 1$, and $\lim_{B\to\infty} \frac{\log(1-\gamma_B)}{B^2} = -\frac{1}{2}$ both follow from Theorem 5, Lemma 27, and Lemma 17.i. ☐

8. Explicit Bounds on the Distance to Stationarity. In this section we prove Theorem 7 by demonstrating some novel bounds on the $\{Q_{n,k}(x)\}$ polynomials, and combining with Theorem 9 and Corollary 1. We begin by bounding $|Q_{n,n}(x)|$, $|Q_{n,n-1}(x)|$, and $|Q_{n,n}(x) - Q_{n,n-1}(x)|$ in terms of $c_n(x)$ and $b_n(x)$. Let $h_n(x) \triangleq 2nb_n(x)^{-1}$.

LEMMA 28. *$x \in \big((\sqrt{n} - \sqrt{\lambda_n})^2, (\sqrt{n} + \sqrt{\lambda_n})^2\big)$ implies*

(i) $|Q_{n,n}(x)| c_n(x)^{-\frac{1}{2}} \leq h_n(x)$,
(ii) $|Q_{n,n-1}(x)| c_n(x)^{-\frac{1}{2}} \leq h_n(x)$,
(iii) $|Q_{n,n}(x) - Q_{n,n-1}(x)| c_n(x)^{-\frac{1}{2}} \leq (\frac{x}{n})^{\frac{1}{2}} h_n(x)$.

PROOF. Expanding $Q_{n,n+1}(x)$ using the recursive definition 12 for the $Q_{n,k}(x)$ polynomials,

$$\begin{aligned}
c_n(x) &= Q^2_{n,n}(x) - Q_{n,n-1}(x)\Big(\frac{\lambda_n + n - x}{\lambda_n} Q_{n,n}(x) - \frac{n}{\lambda_n} Q_{n,n-1}(x)\Big) \\
&= Q^2_{n,n}(x) - \frac{\lambda_n + n - x}{\lambda_n} Q_{n,n}(x) Q_{n,n-1}(x) + \frac{n}{\lambda_n} Q^2_{n,n-1}(x). \qquad (86)
\end{aligned}$$

We now prove assertion i. First, note that $b_n(x) > 0$ and $c_n(x) = Q^2_{n,n}(x) - Q_{n,n-1}(x) Q_{n,n+1}(x) > 0$ by (11) and (13), ensuring that $c_n(x)^{-1}$ and $b_n(x)^{-1}$ are well-defined. If $Q_{n,n}(x) = 0$, then $|Q_{n,n}(x)| c_n(x)^{-\frac{1}{2}} = 0 < h_n(x)$. If $Q_{n,n}(x) \neq 0$ then

$$\begin{aligned}
Q^2_{n,n}(x) c_n(x)^{-1} &= \Big(1 - \frac{\lambda_n + n - x}{\lambda_n} \frac{Q_{n,n-1}(x)}{Q_{n,n}(x)} + \frac{n}{\lambda_n} \big(\frac{Q_{n,n-1}(x)}{Q_{n,n}(x)}\big)^2\Big)^{-1} \\
&\leq \sup_z \Big(1 - \frac{\lambda_n + n - x}{\lambda_n} z + \frac{n}{\lambda_n} z^2\Big)^{-1}. \qquad (87)
\end{aligned}$$

By elementary calculus, $\alpha_1(z) \triangleq 1 - \frac{\lambda_n + n - x}{\lambda_n} z + \frac{n}{\lambda_n} z^2$ is a convex function of z minimized at

$z = \frac{\lambda_n+n-x}{2n}$, and

$$\begin{aligned} \alpha_1(\frac{\lambda_n+n-x}{2n}) &= 1 - \frac{\lambda_n+n-x}{\lambda_n}\frac{\lambda_n+n-x}{2n} + \frac{n}{\lambda_n}(\frac{\lambda_n+n-x}{2n})^2 \\ &= 1 - \frac{(\lambda_n+n-x)^2}{2\lambda_n n} + \frac{(\lambda_n+n-x)^2}{4\lambda_n n} \\ &= \frac{4\lambda_n n - (\lambda_n+n-x)^2}{4\lambda_n n} \quad = \quad \frac{b_n^2(x)}{4\lambda_n n}. \end{aligned}$$

Combining with (87), taking the square root of both sides, and observing that $2\sqrt{\lambda_n n} < 2n$ completes the proof of i.

We now prove assertion ii. If $Q_{n,n-1}(x) = 0$, then $|Q_{n,n-1}(x)|c_n(x)^{-\frac{1}{2}} = 0 < h_n(x)$. If $Q_{n,n-1}(x) \neq 0$ then

$$\begin{aligned} Q_{n,n-1}^2(x)c_n(x)^{-1} &= ((\frac{Q_{n,n}(x)}{Q_{n,n-1}(x)})^2 - \frac{\lambda_n+n-x}{\lambda_n}\frac{Q_{n,n}(x)}{Q_{n,n-1}(x)} + \frac{n}{\lambda_n})^{-1} \\ &\leq \sup_z \left(z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n}\right)^{-1}. \end{aligned} \tag{88}$$

Letting $\alpha_2(z) \overset{\Delta}{=} z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n}$, we find by elementary calculus that $\alpha_2(z)$ is a convex function of z minimized at $z = \frac{\lambda_n+n-x}{2\lambda_n}$, and

$$\begin{aligned} \alpha_2(\frac{\lambda_n+n-x}{2\lambda_n}) &= (\frac{\lambda_n+n-x}{2\lambda_n})^2 - \frac{\lambda_n+n-x}{\lambda_n}\frac{\lambda_n+n-x}{2\lambda_n} + \frac{n}{\lambda_n} \\ &= \frac{(\lambda_n+n-x)^2}{4\lambda_n^2} - \frac{(\lambda_n+n-x)^2}{2\lambda_n^2} + \frac{n}{\lambda_n} \\ &= \frac{4\lambda_n n - (\lambda_n+n-x)^2}{4\lambda_n^2} \quad = \quad \frac{b_n^2(x)}{4\lambda_n^2}. \end{aligned}$$

Combining with (88), taking the square root of both sides, and observing that $2\lambda_n < 2n$ completes the proof of ii.

We now prove assertion iii. It is shown in [38] that $Q_{n,n}(x)$ and $Q_{n,n-1}(x)$ do not have any common zeros. Thus first suppose $Q_{n,n}(x) = 0$. Then from (86), $c_n(x) = \frac{n}{\lambda_n}Q_{n,n-1}^2(x)$. Thus

$$\big(Q_{n,n}(x) - Q_{n,n-1}(x)\big)^2 c_n(x)^{-1} = \frac{Q_{n,n-1}^2(x)}{\frac{n}{\lambda_n}Q_{n,n-1}^2(x)} = \frac{\lambda_n}{n}. \tag{89}$$

Furthermore,

$$\begin{aligned} \frac{4\lambda_n x}{b_n^2(x)} &= \frac{4\lambda_n x}{4\lambda_n n - (\lambda_n+n-x)^2} \\ &= 1 + \frac{4\lambda_n x - 4\lambda_n n + (\lambda_n+n-x)^2}{b_n^2(x)} \\ &= 1 + \frac{4\lambda_n(x-n) + \lambda_n^2 - 2\lambda_n(x-n) + (x-n)^2}{b_n^2(x)} \\ &= 1 + \frac{(\lambda_n+x-n)^2}{b_n^2(x)} \quad \geq \quad 1. \end{aligned} \tag{90}$$

Combining (89) and (90) with the fact that $\frac{\lambda_n}{n} < 1$, we find that

$$\big(Q_{n,n}(x) - Q_{n,n-1}(x)\big)^2 c_n(x)^{-1} \leq \frac{4\lambda_n x}{b_n^2(x)}. \tag{91}$$

Now suppose $Q_{n,n-1}(x) = 0$. Then from (86), $c_n(x) = Q^2_{n,n}(x)$. Thus

$$\big(Q_{n,n}(x) - Q_{n,n-1}(x)\big)^2 c_n(x)^{-1} \;=\; \frac{Q^2_{n,n}(x)}{Q^2_{n,n}(x)} \;=\; 1 \;\leq\; \frac{4\lambda_n x}{b_n^2(x)} \text{ by (90).} \tag{92}$$

Now suppose $Q_{n,n}(x) \neq 0$ and $Q_{n,n-1}(x) \neq 0$. Then

$$\begin{aligned}\big(Q_{n,n}(x) - Q_{n,n-1}(x)\big)^2 c_n(x)^{-1} \;&=\; \frac{(\frac{Q_{n,n}(x)}{Q_{n,n-1}(x)} - 1)^2}{(\frac{Q_{n,n}(x)}{Q_{n,n-1}(x)})^2 - \frac{\lambda_n+n-x}{\lambda_n}\frac{Q_{n,n}(x)}{Q_{n,n-1}(x)} + \frac{n}{\lambda_n}} \\ &\leq\; \sup_z \frac{(z-1)^2}{z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n}}.\end{aligned} \tag{93}$$

Let $\alpha_3(z) \overset{\Delta}{=} \frac{(z-1)^2}{z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n}}$. $\inf_{z\in\mathbb{R}}(z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n}) > 0$ from our earlier analysis of $\alpha_2(z)$. Thus by elementary calculus $\alpha_3(z)$ is a continuously differentiable rational function of z on $\mathbb{R}$, and

$$\begin{aligned}\frac{d}{dz}\alpha_3(z) \;&=\; \frac{2(z-1)(z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n}) - (z-1)^2(2z - \frac{\lambda_n+n-x}{\lambda_n})}{(z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n})^2} \\ &=\; \frac{(z-1)\big(2z^2 - 2\frac{\lambda_n+n-x}{\lambda_n}z + 2\frac{n}{\lambda_n} - 2z(z-1) + \frac{\lambda_n+n-x}{\lambda_n}(z-1)\big)}{(z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n})^2} \\ &=\; \frac{(z-1)\big((n-\lambda_n+x) + (\lambda_n - n + x)z\big)}{\lambda_n(z^2 - \frac{\lambda_n+n-x}{\lambda_n}z + \frac{n}{\lambda_n})^2}.\end{aligned}$$

It follows that the zeros of $\frac{d}{dz}\alpha_3(z)$ occur at $z = 1$ and $z = \frac{\lambda_n-n-x}{\lambda_n-n+x}$. Thus by elementary calculus, $\sup_z \alpha_3(z)$ must be one of $\alpha_3(1), \alpha_3(\frac{\lambda_n-n-x}{\lambda_n-n+x}), \lim_{z\to-\infty}\alpha_3(z), \lim_{z\to\infty}\alpha_3(z)$. Trivially, $\alpha_3(1) = 0$, and $\lim_{z\to-\infty}\alpha_3(z) = \lim_{z\to\infty}\alpha_3(z) = 1$. Furthermore,

$$
\begin{aligned}
\alpha_3\Big(\frac{\lambda_n - n - x}{\lambda_n - n + x}\Big) &= \alpha_3\Big(1 - \frac{2x}{\lambda_n - n + x}\Big) \\
&= \frac{(\frac{2x}{\lambda_n - n + x})^2}{(1 - \frac{2x}{\lambda_n - n + x})^2 - \frac{\lambda_n + n - x}{\lambda_n}(1 - \frac{2x}{\lambda_n - n + x}) + \frac{n}{\lambda_n}} \\
&= \frac{(\frac{2x}{\lambda_n - n + x})^2}{1 - \frac{4x}{\lambda_n - n + x} + (\frac{2x}{\lambda_n - n + x})^2 - \frac{\lambda_n + n - x}{\lambda_n} + \frac{\lambda_n + n - x}{\lambda_n - n + x}\frac{2x}{\lambda_n} + \frac{n}{\lambda_n}} \\
&= \frac{(\frac{2x}{\lambda_n - n + x})^2}{1 - \frac{4x}{\lambda_n - n + x} + (\frac{2x}{\lambda_n - n + x})^2 - \frac{\lambda_n + n - x}{\lambda_n} + (1 + \frac{2(n-x)}{\lambda_n - n + x})\frac{2x}{\lambda_n} + \frac{n}{\lambda_n}} \\
&= \frac{\frac{4x}{(\lambda_n - n + x)^2}}{\frac{-4}{\lambda_n - n + x} + \frac{4x}{(\lambda_n - n + x)^2} + \frac{3}{\lambda_n} + \frac{4n - 4x}{\lambda_n(\lambda_n - n + x)}} \\
&= \frac{4x\lambda_n}{-4\lambda_n(\lambda_n - n + x) + 4x\lambda_n + 4(n - x)(\lambda_n - n + x) + 3(\lambda_n - n + x)^2} \\
&= \frac{4x\lambda_n}{(\lambda_n - n + x)\big(-4\lambda_n + 4(n - x) + 3(\lambda_n - n + x)\big) + 4x\lambda_n} \\
&= \frac{4x\lambda_n}{(\lambda_n - n + x)(-\lambda_n + n - x) + 4x\lambda_n} \\
&= \frac{4x\lambda_n}{4x\lambda_n - (\lambda_n - n + x)^2} \quad = \quad \frac{4x\lambda_n}{b_n^2(x)}. \qquad (94)
\end{aligned}
$$

Combining the above with (90), we find that $\sup_z \alpha_3(z) \le \frac{4\lambda_n x}{b_n^2(x)}$.

$$
\begin{aligned}
\sup_z \alpha_3(z) &\le \max\Big(\alpha_3(1), \alpha_3\Big(\frac{\lambda_n - n - x}{\lambda_n - n + x}\Big), \lim_{z\to-\infty}\alpha_3(z), \lim_{z\to\infty}\alpha_3(z)\Big) \\
&= \max\Big(0, \frac{4\lambda_n x}{b_n^2(x)}, 1, 1\Big) \\
&= \frac{4\lambda_n x}{b_n^2(x)} \text{ by (90).} \qquad (95)
\end{aligned}
$$

Combining with (91) - (93) and taking square roots completes the proof of the lemma. □

We now prove some simple asymptotic bounds that are necessary to proceed with our analysis. Recall that $q_n = 1 + n^{-\frac{1}{4}}$. Let $r_n(x) \overset{\Delta}{=} q_n x n^{-1}$, and $s_n(\alpha) \overset{\Delta}{=} q_n \alpha n^{-\frac{1}{2}}$. Then

LEMMA 29. *For each $\alpha \ge B > 0$, there exists N_α s.t. $n \ge N_\alpha$, $k \in [n - \alpha\sqrt{n} - 3, n + \alpha\sqrt{n} + 3]$, and $x > 0$ implies*

(i) $|\frac{x}{\min(k,n)}|, |\frac{x}{\lambda_n}| \le r_n(x)$,
(ii) $|\frac{\lambda_n}{\min(k,n)}|, |\frac{n}{\lambda_n}| \le 1 + s_n(\alpha)$.

PROOF. For any constants $a, b > 0$ and all sufficiently large n,

$$\begin{aligned}\frac{n}{n - an^{\frac{1}{2}} - b} &= 1 + \frac{an^{\frac{1}{2}} + b}{n - an^{\frac{1}{2}} - b} \\ &= 1 + an^{-\frac{1}{2}} + \frac{(a^2 + b)n + abn^{\frac{1}{2}}}{n^2 - an^{\frac{3}{2}} - bn} \\ &= 1 + an^{-\frac{1}{2}} + O(n^{-1}) \\ &\leq 1 + an^{-\frac{1}{2}} + an^{-\frac{3}{4}} \text{ for all sufficiently large n} \\ &= 1 + s_n(a). \qquad (96)\end{aligned}$$

Similarly, for any constants $a, b > 0$ and all sufficiently large n,

$$\begin{aligned}(n - an^{\frac{1}{2}} - b)^{-1} &\leq \big(1 + s_n(a)\big)n^{-1} \text{ by (96)} \\ &= n^{-1} + q_n an^{-\frac{3}{2}} \\ &\leq n^{-1} + n^{-\frac{5}{4}} \text{ for all sufficiently large n, since } \ q_n an^{-\frac{3}{2}} = O(n^{-\frac{3}{2}}) \\ &= q_n n^{-1}. \qquad (97)\end{aligned}$$

Thus from the assumptions on k, the definition of λ_n, and the assumption that $\alpha \geq B$, we find that for all sufficiently large n,

$$\min(k, n)^{-1} \leq q_n n^{-1} \text{ by taking } \ a = \alpha, b = 3 \text{ in (97)}; \qquad (98)$$

$$\lambda_n^{-1} \leq q_n n^{-1} \text{ by taking } a = \alpha, b = 3 \text{ in (97) since } \lambda_n^{-1} \leq (n - \alpha n^{\frac{1}{2}} - 3)^{-1}; \qquad (99)$$

$$\frac{\lambda_n}{\min(k - 1, n)} \leq 1 + s_n(\alpha) \text{ since } \lambda_n \leq n, \text{ and applying (96) with } \ a = \alpha, b = 4; \qquad (100)$$

$$\frac{n}{\lambda_n} \leq 1 + s_n(\alpha) \text{ by applying (96) with } \ a = \alpha, b = 0. \qquad (101)$$

Assertions i. and ii. follow immediately from (98)-(101) and the assumed positivity of x. ☐

We now bound $|Q_{n,k+1}(x)|$ and $|Q_{n,k+1}(x) - Q_{n,k}(x)|$ in terms of $|Q_{n,k}(x)|$ and $|Q_{n,k}(x) - Q_{n,k-1}(x)|$ and visa-versa, which will later enable us to bound $|Q_{n,k}(x)|$ for $k = n \pm O(\sqrt{n})$ using Lemma 28. More precisely, we prove

LEMMA 30. *For each $\alpha \geq B > 0$, there exists N_α s.t. $n \geq N_\alpha$, $k \in [n - \alpha\sqrt{n} - 3, n + \alpha\sqrt{n} + 3]$, and $x > 0$ implies*

(i) $|Q_{n,k+1}(x)| \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(|Q_{n,k}(x)| + |Q_{n,k}(x) - Q_{n,k-1}(x)|\big)$,
(ii) $|Q_{n,k+1}(x) - Q_{n,k}(x)| \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(r_n(x)|Q_{n,k}(x)| + |Q_{n,k}(x) - Q_{n,k-1}(x)|\big)$,
(iii) $|Q_{n,k-1}(x)| \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(|Q_{n,k}(x)| + |Q_{n,k}(x) - Q_{n,k+1}(x)|\big)$,
(iv) $|Q_{n,k-1}(x) - Q_{n,k}(x)| \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(r_n(x)|Q_{n,k}(x)| + |Q_{n,k}(x) - Q_{n,k+1}(x)|\big)$.

Proof. Let us choose N_α to satisfy the conditions of Lemma 29 (the existence of such an N_α follows from the same lemma), and suppose $n \geq N_\alpha$. We first prove assertion i. Recall that by for $k \geq 1$, $|Q_{n,k+1}(x)|$ is equivalent to

$$|(1 - \frac{x}{\lambda_n} + \frac{\min(k,n)}{\lambda_n})Q_{n,k}(x) - \frac{\min(k,n)}{\lambda_n}Q_{n,k-1}(x)|$$

$$\begin{aligned}
&= |(1 - \frac{x}{\lambda_n})Q_{n,k}(x) + \frac{\min(k,n)}{\lambda_n}\big(Q_{n,k}(x) - Q_{n,k-1}(x)\big)| \\
&\leq |1 - \frac{x}{\lambda_n}||Q_{n,k}(x)| + |\frac{\min(k,n)}{\lambda_n}||Q_{n,k}(x) - Q_{n,k-1}(x)| \quad \text{by the triangle inequality} \\
&\leq \big(1 + r_n(x)\big)|Q_{n,k}(x)| + \big(1 + s_n(\alpha)\big)|Q_{n,k}(x) - Q_{n,k-1}(x)| \quad \text{by Lemma 29.i and Lemma 29.ii,}
\end{aligned}$$

from which assertion i. follows, since by the strict positivity of $r_n(x)$ and $s_n(\alpha)$ we have

$$\big(1 + r_n(x)\big) \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big), \text{ and } \big(1 + s_n(\alpha)\big) \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big). \tag{102}$$

We now prove assertion ii. Rearranging definition (12), we find that for $k \geq 1$, $|Q_{n,k+1}(x) - Q_{n,k}(x)|$ is equivalent to

$$|-\frac{x}{\lambda_n}Q_{n,k}(x) + \frac{\min(k,n)}{\lambda_n}\big(Q_{n,k}(x) - Q_{n,k-1}(x)\big)|$$

$$\begin{aligned}
&\leq |-\frac{x}{\lambda_n}||Q_{n,k}(x)| + |\frac{\min(k,n)}{\lambda_n}||Q_{n,k}(x) - Q_{n,k-1}(x)| \quad \text{by the triangle inequality} \\
&\leq r_n(x)|Q_{n,k}(x)| + \big(1 + s_n(\alpha)\big)|Q_{n,k}(x) - Q_{n,k-1}(x)| \quad \text{by Lemma 29.i and Lemma 29.ii,}
\end{aligned}$$

from which assertion ii. follows by (102) and the similar fact that

$$r_n(x) \leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)r_n(x). \tag{103}$$

We now prove assertion iii. Rearranging definition (12), we find that for $k \geq 1$, $|Q_{n,k-1}(x)|$ is equivalent to

$$|\frac{\lambda_n}{\min(k,n)}\big((1 - \frac{x}{\lambda_n} + \frac{\min(k,n)}{\lambda_n})Q_{n,k}(x) - Q_{n,k+1}(x)\big)|$$

$$\begin{aligned}
&= |(\frac{\lambda_n}{\min(k,n)} - \frac{x}{\min(k,n)} + 1)Q_{n,k}(x) - \frac{\lambda_n}{\min(k,n)}Q_{n,k+1}(x)| \\
&= |\big(1 - \frac{x}{\min(k,n)}\big)Q_{n,k}(x) + \frac{\lambda_n}{\min(k,n)}\big(Q_{n,k}(x) - Q_{n,k+1}(x)\big)| \\
&\leq |1 - \frac{x}{\min(k,n)}||Q_{n,k}(x)| + |\frac{\lambda_n}{\min(k,n)}||Q_{n,k}(x) - Q_{n,k+1}(x)| \quad \text{by the triangle inequality} \\
&\leq \big(1 + r_n(x)\big)|Q_{n,k}(x)| + \big(1 + s_n(\alpha)\big)|Q_{n,k}(x) - Q_{n,k+1}(x)| \quad \text{by Lemma 29.i and Lemma 29.ii,}
\end{aligned}$$

from which assertion iii. follows by (102).

We now prove assertion iv. Rearranging definition (12), we find that for $k \geq 1$, $|Q_{n,k-1}(x) - Q_{n,k}(x)|$ is equivalent to

$$|\frac{\lambda_n}{\min(k,n)}\big((1 - \frac{x}{\lambda_n} + \frac{\min(k,n)}{\lambda_n})Q_{n,k}(x) - Q_{n,k+1}(x)\big) - Q_{n,k}(x)|$$

$$
\begin{aligned}
&= |(\frac{\lambda_n}{\min(k,n)} - \frac{x}{\min(k,n)})Q_{n,k}(x) - \frac{\lambda_n}{\min(k,n)}Q_{n,k+1}(x)| \\
&= |-\frac{x}{\min(k,n)}Q_{n,k}(x) + \frac{\lambda_n}{\min(k,n)}\big(Q_{n,k}(x) - Q_{n,k+1}(x)\big)| \\
&\leq |-\frac{x}{\min(k,n)}||Q_{n,k}(x)| + |\frac{\lambda_n}{\min(k,n)}||Q_{n,k}(x) - Q_{n,k+1}(x)| \quad \text{by the triangle inequality} \\
&\leq r_n(x)|Q_{n,k}(x)| + \big(1 + s_n(\alpha)\big)|Q_{n,k}(x) - Q_{n,k+1}(x)| \quad \text{by Lemma 29.i and Lemma 29.ii,}
\end{aligned}
$$

from which assertion iv. follows by (102) and (103). ☐

We now use Lemma 30 to inductively bound $|Q_{n,k}(x)|$ for values of k that are $O(n^{\frac{1}{2}})$ away from n.

LEMMA 31. *For each $\alpha \geq B > 0$, there exists N_α s.t. $n \geq N_\alpha$, and $x \in \big((\sqrt{n} - \sqrt{\lambda_n})^2, (\sqrt{n} + \sqrt{\lambda_n})^2\big)$ implies that*

(i) For $k \in [n, n + \alpha\sqrt{n} + 3]$,

  (a) $|Q_{n,k}(x)| \leq \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{k-n} c_n(x)^{\frac{1}{2}} h_n(x)$,

  (b) $|Q_{n,k}(x) - Q_{n,k-1}(x)| \leq \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{k-n} r_n(x)^{\frac{1}{2}} c_n(x)^{\frac{1}{2}} h_n(x)$.

(ii) For $k \in [n - \alpha\sqrt{n} - 3, n - 1]$,

  (a) $|Q_{n,k}(x)| \leq \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{n-k} c_n(x)^{\frac{1}{2}} h_n(x)$,

  (b) $|Q_{n,k}(x) - Q_{n,k+1}(x)| \leq \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{n-k} r_n(x)^{\frac{1}{2}} c_n(x)^{\frac{1}{2}} h_n(x)$.

PROOF. Let us choose N_α to satisfy the conditions of Lemma 30 (the existence of such an N_α follows from the same lemma), and suppose $n \geq N_\alpha$. We first prove assertion i. We proceed by induction on ia. and ib. simultaneously. The base case $k = n$ follows immediately from Lemma 28.i and Lemma 28.iii. Now, suppose the induction is true for some $k \in [n, n + \alpha n^{\frac{1}{2}} + 2]$. Then by Lemma 30.i,

$$
\begin{aligned}
|Q_{n,k+1}(x)| &\leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(|Q_{n,k}(x)| + |Q_{n,k}(x) - Q_{n,k-1}(x)|\big) \\
&\leq \big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\Bigg(\Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{k-n} c_n(x)^{\frac{1}{2}} h_n(x) \\
&\quad + \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{k-n} r_n(x)^{\frac{1}{2}} c_n(x)^{\frac{1}{2}} h_n(x)\Bigg) \quad \text{by the ind. hyp.} \\
&= \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\Big)^{k+1-n}\big(1 + r_n(x)^{\frac{1}{2}}\big)^{k-n} c_n(x)^{\frac{1}{2}} h_n(x)\big(1 + r_n(x)^{\frac{1}{2}}\big) \\
&= \Big(\big(1 + r_n(x)\big)\big(1 + s_n(\alpha)\big)\big(1 + r_n(x)^{\frac{1}{2}}\big)\Big)^{k+1-n} c_n(x)^{\frac{1}{2}} h_n(x). \qquad (104)
\end{aligned}
$$

Similarly, by Lemma 30.ii

$$\begin{aligned}
|Q_{n,k+1}(x)-Q_{n,k}(x)| &\le \big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(r_n(x)|Q_{n,k}(x)|+|Q_{n,k}(x)-Q_{n,k-1}(x)|\big)\\
&\le \big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\Bigg(\Big(\big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(1+r_n(x)^{\frac{1}{2}}\big)\Big)^{k-n} r_n(x)c_n(x)^{\frac{1}{2}}h_n(x)\\
&\quad + \Big(\big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(1+r_n(x)^{\frac{1}{2}}\big)\Big)^{k-n} r_n(x)^{\frac{1}{2}}c_n(x)^{\frac{1}{2}}h_n(x)\Bigg) \quad \text{by the ind. hyp.}\\
&= \Big(\big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\Big)^{k+1-n}\big(1+r_n(x)^{\frac{1}{2}}\big)^{k-n}c_n(x)^{\frac{1}{2}}h_n(x)\big(r_n(x)+r_n(x)^{\frac{1}{2}}\big)\\
&= \Big(\big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(1+r_n(x)^{\frac{1}{2}}\big)\Big)^{k+1-n} r_n(x)^{\frac{1}{2}}c_n(x)^{\frac{1}{2}}h_n(x). \qquad (105)
\end{aligned}$$

This concludes the induction, proving assertion i.

We now prove assertion ii., which follows from a very similar argument. We proceed by induction on iia. and iib. simultaneously. The base case $k=n-1$ follows immediately from Lemma 28.ii and Lemma 28.iii. Now, suppose the induction is true for some $k\in[n-\alpha n^{\frac{1}{2}}-2, n-1]$. Then by Lemma 30.iii,

$$\begin{aligned}
|Q_{n,k-1}(x)| &\le \big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(|Q_{n,k}(x)|+|Q_{n,k}(x)-Q_{n,k+1}(x)|\big)\\
&\le \Big(\big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(1+r_n(x)^{\frac{1}{2}}\big)\Big)^{n-(k-1)} c_n(x)^{\frac{1}{2}}h_n(x)\\
&\quad \text{by exactly the same argument used to derive (104).}
\end{aligned}$$

Similarly, by Lemma 30.iv,

$$\begin{aligned}
|Q_{n,k-1}(x)-Q_{n,k}(x)| &\le \big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(r_n(x)|Q_{n,k}(x)|+|Q_{n,k}(x)-Q_{n,k+1}(x)|\big)\\
&\le \Big(\big(1+r_n(x)\big)\big(1+s_n(\alpha)\big)\big(1+r_n(x)^{\frac{1}{2}}\big)\Big)^{n-k} r_n(x)^{\frac{1}{2}}c_n(x)^{\frac{1}{2}}h_n(x)\\
&\quad \text{by exactly the same argument used to derive (105).}
\end{aligned}$$

This concludes the induction, proving assertion ii. and thus the lemma. □

LEMMA 32. *For each $\alpha\ge B>0$, there exists N_α s.t. $n\ge N_\alpha$, $x\in\big((\sqrt{n}-\sqrt{\lambda_n})^2,(\sqrt{n}+\sqrt{\lambda_n})^2\big)$, and $k\in[n-\alpha\sqrt{n}-2, n+\alpha\sqrt{n}+2]$ implies that*

(i) $|Q_{n,k}(x)|c_n(x)^{-\frac{1}{2}}\le h_n(x)\exp\big((\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)$

(ii) $|Q_{n,k+1}(x)-Q_{n,k}(x)|c_n(x)^{-\frac{1}{2}}\le h_n(x)r_n(x)^{\frac{1}{2}}\exp\big((\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)$

PROOF. By Lemma 31.ia and Lemma 31.iia, we find that there exists N_α s.t. $n\ge N_\alpha$ implies

$$\begin{aligned}
|Q_{n,k}(x)|c_n(x)^{-\frac{1}{2}} &\le h_n(x)\exp\big(|n-k|(r_n(x)+s_n(\alpha)+r_n(x)^{\frac{1}{2}})\big)\\
&\le h_n(x)\exp\big((\alpha n^{\frac{1}{2}}+2)(r_n(x)+s_n(\alpha)+r_n(x)^{\frac{1}{2}})\big)\\
&= h_n(x)\exp\Big(\big(\alpha n^{\frac{1}{2}}+2\big)\big(q_n x n^{-1}+q_n\alpha n^{-\frac{1}{2}}+q_n^{\frac{1}{2}}x^{\frac{1}{2}}n^{-\frac{1}{2}}\big)\Big)\\
&\le h_n(x)\exp\Big(q_n\big(\alpha n^{\frac{1}{2}}+2\big)\big(xn^{-1}+\alpha n^{-\frac{1}{2}}+x^{\frac{1}{2}}n^{-\frac{1}{2}}\big)\Big) \quad \text{since} \quad q_n^{\frac{1}{2}}\le q_n\\
&= h_n(x)\exp\Big(q_n\big(\alpha+2n^{-\frac{1}{2}}\big)\big(xn^{-\frac{1}{2}}+\alpha+x^{\frac{1}{2}}\big)\Big). \qquad (106)
\end{aligned}$$

By identical reasoning,

$$|Q_{n,k+1}(x)-Q_{n,k}(x)|c_n(x)^{-\frac{1}{2}} \le h_n(x)r_n(x)^{\frac{1}{2}}\exp\Big(q_n\big(\alpha+2n^{-\frac{1}{2}}\big)\big(xn^{-\frac{1}{2}}+\alpha+x^{\frac{1}{2}}\big)\Big). \tag{107}$$

Furthermore, note that since

$$\frac{xn^{-\frac{1}{2}}}{2x^{\frac{1}{2}}} = \frac{x^{\frac{1}{2}}}{2n^{\frac{1}{2}}} < \frac{n^{\frac{1}{2}}+\lambda_n^{\frac{1}{2}}}{2n^{\frac{1}{2}}} < 1, \tag{108}$$

it will be the case that

$$0 < x < (\sqrt{n}+\sqrt{\lambda_n})^2 \text{ implies } xn^{-\frac{1}{2}} < 2x^{\frac{1}{2}}. \tag{109}$$

Also, note that

$$\begin{aligned} q_n\big(\alpha+2n^{-\frac{1}{2}}\big) &= (1+n^{-\frac{1}{4}})(\alpha+2n^{-\frac{1}{2}}) \\ &= \alpha+\alpha n^{-\frac{1}{4}}+2n^{-\frac{1}{2}}+2n^{-\frac{3}{4}} \\ &= \alpha+O(n^{-\frac{1}{4}}) \\ &\le \alpha+n^{-\frac{1}{5}} \quad \text{for all sufficiently large n.} \qquad (110) \end{aligned}$$

The lemma follows by combining (106), (107),(109), and (110). ☐

We now derive a variant of Theorem 9 for $P^n_{i,\le j}(t)$ (as opposed to $P^n_{i,j}(t)$) that does not simply sum over all $j+1$ states $\le j$, but instead relies on a ‘probability flow’ interpretation using the Chapman-Kolmogorov equation. This will allow us to bound the distance to stationarity uniformly in n for the c.d.f., since we can sidestep having to sum over ‘too many’ error terms.

LEMMA 33. *For* $\frac{\lambda_n}{n} \ge \rho^*_n$,

$$|P^n_{i,\le j}(t)-P^n_{\le j}(\infty)| \le (2\pi)^{-1}g_n(j)\frac{n!}{\lambda_n^n}n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2} e^{-xt}x^{-1}|Q_{n,i}(x)||Q_{n,j+1}(x)-Q_{n,j}(x)|b_n(x)c_n(x)^{-1}dx. \tag{111}$$

PROOF. Note that

$$\begin{aligned} \frac{d}{dt}P^n_{i,\le j}(t) &= \sum_{k=0}^{j}\frac{d}{dt}P^n_{i,k}(t) \\ &= P^n_{i,1}(t)-\lambda_n P^n_{i,0}(t)+\sum_{k=1}^{j}\big(\min(k+1,n)P^n_{i,k+1}(t)+\lambda_n P^n_{i,k-1}(t)-(\lambda_n+\min(k,n))P^n_{i,k}(t)\big) \\ &\qquad \text{by the Chapman-Kolmogorov equation} \\ &= \min(j+1,n)P^n_{i,j+1}(t)-\lambda_n P^n_{i,j}(t) \quad \text{by a telescoping sum argument.} \end{aligned}$$

It follows that for all $i,t\ge 0$, $P^n_{\le j}(\infty)=P^n_{i,\le j}(t)+\int_t^\infty\big(\min(j+1,n)P^n_{i,j+1}(s)-\lambda_n P^n_{i,j}(s)\big)ds$. Thus

$$|P^n_{i,\le j}(t)-P^n_{\le j}(\infty)|$$

$$
\begin{aligned}
&= \Big|\int_t^{\infty}\big(\min(j+1,n)P^n_{i,j+1}(s)-\lambda_n P^n_{i,j}(s)\big)ds\Big| \\
&= \Big|\int_t^{\infty}\Big(\min(j+1,n)\big(P^n_{j+1}(\infty)+(P^n_{i,j+1}(s)-P^n_{i,j+1}(\infty))\big)-\lambda_n\big(P^n_j(\infty)+(P^n_{i,j}(s)-P^n_{i,j}(\infty))\big)\Big)ds\Big| \\
&= \Big|\int_t^{\infty}\Big(\min(j+1,n)\big(P^n_{i,j+1}(s)-P^n_{i,j+1}(\infty)\big)-\lambda_n\big(P^n_{i,j}(s)-P^n_{i,j}(\infty)\big)\Big)ds\Big| \\
&\qquad \text{since the steady-state terms cancel by detailed balance} \\
&= \Bigg|\int_t^{\infty}\Bigg(\min(j+1,n)\Big((2\pi)^{-1}g_n(j+1)\frac{n!}{\lambda_n^n}(\lambda_n n)^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}e^{-xs}Q_{n,i}(x)Q_{n,j+1}(x)b_n(x)c_n(x)^{-1}dx\Big) \\
&\qquad -\lambda_n\Big((2\pi)^{-1}g_n(j)\frac{n!}{\lambda_n^n}(\lambda_n n)^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}e^{-xs}Q_{n,i}(x)Q_{n,j}(x)b_n(x)c_n(x)^{-1}dx\Big)\Bigg)ds\Bigg| \quad \text{by Theorem 9} \\
&= \Big|\int_t^{\infty}\Big((2\pi)^{-1}g_n(j)\frac{n!}{\lambda_n^n}n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}e^{-xs}Q_{n,i}(x)\big(Q_{n,j+1}(x)-Q_{n,j}(x)\big)b_n(x)c_n(x)^{-1}dx\Big)ds\Big| \\
&\qquad \text{since } \min(j+1,n)g_n(j+1)\frac{n!}{\lambda_n^n}(\lambda_n n)^{-1}=\lambda_n g_n(j)\frac{n!}{\lambda_n^n}(\lambda_n n)^{-1}=g_n(j)\frac{n!}{\lambda_n^n}n^{-1} \\
&\leq \int_t^{\infty}\Big((2\pi)^{-1}g_n(j)\frac{n!}{\lambda_n^n}n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}e^{-xs}|Q_{n,i}(x)||Q_{n,j+1}(x)-Q_{n,j}(x)|b_n(x)c_n(x)^{-1}dx\Big)ds \\
&= (2\pi)^{-1}g_n(j)\frac{n!}{\lambda_n^n}n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\int_t^{\infty}e^{-xs}|Q_{n,i}(x)||Q_{n,j+1}(x)-Q_{n,j}(x)|b_n(x)c_n(x)^{-1}ds\ dx \\
&\quad \text{by Tonelli's Theorem} \\
&= (2\pi)^{-1}g_n(j)\frac{n!}{\lambda_n^n}n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}e^{-xt}x^{-1}|Q_{n,i}(x)||Q_{n,j+1}(x)-Q_{n,j}(x)|b_n(x)c_n(x)^{-1}dx,
\end{aligned}
$$

and the lemma follows. ☐

We now prove bounds on a special type of integral that arises in the analysis of $P^n_{i,j}(t)-P^n_j(\infty)$.

LEMMA 34. *For all $B,C>0$ there exists $N_{B,C}$ s.t. $n\geq N_{B,C}$ implies that for all $c\in(0,C]$ and $t\geq 1$,*

$$\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt+cx^{\frac{1}{2}}\big)b_n(x)^{-1}dx\leq q_n\big(\frac{\pi}{t\lambda_n}\big)^{\frac{1}{2}}\exp\big(\frac{1}{2}Bc+c^2(2t)^{-1}-\frac{B^2}{4}t\big).$$

PROOF. By Lemma 4, $(\sqrt{n}-\sqrt{\lambda_n})^2<2\sqrt{\lambda_n n}<(\sqrt{n}+\sqrt{\lambda_n})^2$ for all sufficiently large n. Thus

$$\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt+cx^{\frac{1}{2}}\big)\big(4\lambda_n n-(\lambda_n+n-x)^2\big)^{-\frac{1}{2}}dx$$

$$(112) = \int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{2\sqrt{\lambda_n n}}\exp\big(-xt+cx^{\frac{1}{2}}\big)\big(x-(\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}}\big((\sqrt{n}+\sqrt{\lambda_n})^2-x\big)^{-\frac{1}{2}}dx$$

$$(113)\quad +\int_{2\sqrt{\lambda_n n}}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt+cx^{\frac{1}{2}}\big)\big(x-(\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}}\big((\sqrt{n}+\sqrt{\lambda_n})^2-x\big)^{-\frac{1}{2}}dx \text{ by (11).}$$

Let $u_n \triangleq 2\sqrt{\lambda_n n} - (\sqrt{n}-\sqrt{\lambda_n})^2$. Since $\big((\sqrt{n}+\sqrt{\lambda_n})^2 - x\big)^{-\frac{1}{2}} \le \big((\sqrt{n}+\sqrt{\lambda_n})^2 - 2\sqrt{\lambda_n n}\big)^{-\frac{1}{2}}$ for $x \in \big((\sqrt{n}-\sqrt{\lambda_n})^2, 2\sqrt{\lambda_n n}\big)$, the term appearing in (112) is at most

$$\big((\sqrt{n}+\sqrt{\lambda_n})^2 - 2\sqrt{\lambda_n n}\big)^{-\frac{1}{2}} \int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{2\sqrt{\lambda_n n}} \exp\big(-xt + cx^{\frac{1}{2}}\big)\big(x - (\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}}dx$$

$$\begin{aligned}
&= \big(\lambda_n + n\big)^{-\frac{1}{2}} \int_0^{u_n} \exp\Big(-\big(y + (\sqrt{n}-\sqrt{\lambda_n})^2\big)t + c\big(y + (\sqrt{n}-\sqrt{\lambda_n})^2\big)^{\frac{1}{2}}\Big) y^{-\frac{1}{2}}dy \\
&\qquad \text{substituting } y = x - (\sqrt{n}-\sqrt{\lambda_n})^2 \\
&\le \big(\lambda_n + n\big)^{-\frac{1}{2}} \exp\Big(-\big(\sqrt{n}-\sqrt{\lambda_n})^2\big)t\Big) \int_0^{u_n} \exp\Big(-yt + cy^{\frac{1}{2}} + c(\sqrt{n}-\sqrt{\lambda_n})\Big) y^{-\frac{1}{2}}dy \\
&\qquad \text{since } a, b > 0 \text{ implies } (a+b)^{\frac{1}{2}} < a^{\frac{1}{2}} + b^{\frac{1}{2}} \\
&= \big(\lambda_n + n\big)^{-\frac{1}{2}} \exp\Big(-(\sqrt{n}-\sqrt{\lambda_n})^2 t + c(\sqrt{n}-\sqrt{\lambda_n})\Big) \int_0^{u_n} \exp\big(-yt + cy^{\frac{1}{2}}\big) y^{-\frac{1}{2}}dy.
\end{aligned} \tag{114}$$

We now show that

$$-yt + cy^{\frac{1}{2}} \le -\frac{1}{2}yt + c^2(2t)^{-1} \text{ for all } y > 0. \tag{115}$$

Indeed, by elementary calculus the concave function $\alpha_4(y) \triangleq (-yt + cy^{\frac{1}{2}}) - (-\frac{1}{2}yt + c^2(2t)^{-1})$ attains its maximum when $y = (\frac{c}{t})^2$, and $\alpha_4\big((\frac{c}{t})^2\big) = 0$, from which the desired inequality follows. Plugging (115) into (114), we find that the term appearing in (112) is at most

$$\big(\lambda_n + n\big)^{-\frac{1}{2}} \exp\Big(-(\sqrt{n}-\sqrt{\lambda_n})^2 t + c(\sqrt{n}-\sqrt{\lambda_n})\Big) \int_0^{u_n} \exp\big(-\frac{1}{2}yt + c^2(2t)^{-1}\big) y^{-\frac{1}{2}}dy$$

$$\begin{aligned}
&= \big(\lambda_n + n\big)^{-\frac{1}{2}} \exp\Big(-(\sqrt{n}-\sqrt{\lambda_n})^2 t + c(\sqrt{n}-\sqrt{\lambda_n}) + c^2(2t)^{-1}\Big) \int_0^{u_n} \exp\big(-\frac{1}{2}yt\big) y^{-\frac{1}{2}}dy \\
&\le \big(\lambda_n + n\big)^{-\frac{1}{2}} \exp\Big(-(\sqrt{n}-\sqrt{\lambda_n})^2 t + c(\sqrt{n}-\sqrt{\lambda_n}) + c^2(2t)^{-1}\Big) \int_0^{\infty} \exp\big(-\frac{1}{2}yt\big) y^{-\frac{1}{2}}dy \\
&= \big(\lambda_n + n\big)^{-\frac{1}{2}} \exp\Big(-(\sqrt{n}-\sqrt{\lambda_n})^2 t + c(\sqrt{n}-\sqrt{\lambda_n}) + c^2(2t)^{-1}\Big) (\frac{2\pi}{t})^{\frac{1}{2}} \\
&\qquad \text{since } \int_0^{\infty} \exp\big(-\frac{1}{2}yt\big) y^{-\frac{1}{2}}dy = (\frac{2\pi}{t})^{\frac{1}{2}} \text{ by standard integration techniques} \\
&\le (\frac{\pi}{\lambda_n t})^{\frac{1}{2}} \exp\Big(-(\sqrt{n}-\sqrt{\lambda_n})^2 t + c(\sqrt{n}-\sqrt{\lambda_n}) + c^2(2t)^{-1}\Big) \quad \text{since } \lambda_n + n > 2\lambda_n.
\end{aligned} \tag{116}$$

We now bound the term appearing in (113). Since $\big(x - (\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}} \le \big(2\sqrt{\lambda_n n} - (\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}}$ for $x \in \big(2\sqrt{\lambda_n n}, (\sqrt{n}+\sqrt{\lambda_n})^2\big)$, the term appearing in (113) is at most

$$\big(2\sqrt{\lambda_n n} - (\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}} \int_{2\sqrt{\lambda_n n}}^{(\sqrt{n}+\sqrt{\lambda_n})^2} \exp\big(-xt + cx^{\frac{1}{2}}\big)\big((\sqrt{n}+\sqrt{\lambda_n})^2 - x\big)^{-\frac{1}{2}}dx$$

$$\le \big(2\sqrt{\lambda_n n} - (\sqrt{n}-\sqrt{\lambda_n})^2\big)^{-\frac{1}{2}} \sup_{z \in [2\sqrt{\lambda_n n}, (\sqrt{n}+\sqrt{\lambda_n})^2]} \exp\big(-zt + cz^{\frac{1}{2}}\big) \int_{2\sqrt{\lambda_n n}}^{(\sqrt{n}+\sqrt{\lambda_n})^2} \big((\sqrt{n}+\sqrt{\lambda_n})^2 - x\big)^{-\frac{1}{2}}dx$$

$$= \left(2\sqrt{\lambda_n n} - (\sqrt{n} - \sqrt{\lambda_n})^2\right)^{-\frac{1}{2}} \sup_{z\in[2\sqrt{\lambda_n n},(\sqrt{n}+\sqrt{\lambda_n})^2]} \exp\left(-zt + cz^{\frac{1}{2}}\right) \int_0^{\lambda_n+n} y^{-\frac{1}{2}}dy$$

substituting $y = (\sqrt{n}+\sqrt{\lambda_n})^2 - x$

$$\leq 2\left(\frac{\lambda_n+n}{3\lambda_n-n}\right)^{\frac{1}{2}} \sup_{z\in[2\sqrt{\lambda_n n},(\sqrt{n}+\sqrt{\lambda_n})^2]} \exp\left(-zt + cz^{\frac{1}{2}}\right)$$

since $2\sqrt{\lambda_n n} - (\sqrt{n}-\sqrt{\lambda_n})^2 \geq 3\lambda_n - n,$ and $\int_0^{\lambda_n+n} y^{-\frac{1}{2}}dy = 2(\lambda_n+n)^{\frac{1}{2}}$

$$\leq 2\left(\frac{\lambda_n+n}{3\lambda_n-n}\right)^{\frac{1}{2}} \exp\left(c^2(2t)^{-1} - (\lambda_n n)^{\frac{1}{2}}t\right) \tag{117}$$

by applying (115) to $\sup_{z\in[2\sqrt{\lambda_n n},(\sqrt{n}+\sqrt{\lambda_n})^2]} \exp\left(-zt + cz^{\frac{1}{2}}\right)$.

Furthermore,

$$\left(\frac{\lambda_n+n}{3\lambda_n-n}\right)^{\frac{1}{2}} \leq 2 \text{ for all sufficiently large n, since } \lim_{n\to\infty}\frac{\lambda_n+n}{3\lambda_n-n} = 1. \tag{118}$$

Also,

$$\begin{aligned} c(\sqrt{n}-\sqrt{\lambda_n}) &\leq \frac{1}{2}Bc + \frac{B^2c}{4n^{\frac{1}{2}}-2B} \text{ by Lemma 4} \\ &= \frac{1}{2}Bc + O(n^{-\frac{1}{2}}) \\ &\leq \frac{1}{2}Bc + n^{-\frac{1}{3}} \text{ for all sufficiently large n.} \end{aligned} \tag{119}$$

In addition,

$$\begin{aligned} (\lambda_n n)^{\frac{1}{2}} &\geq \lambda_n \\ &\geq \frac{1}{2}\lambda_n + \lambda_n^{\frac{1}{2}} + \frac{B^2}{4} \text{ for all sufficiently large n,} \\ &\quad \text{since } \lambda_n^{\frac{1}{2}} + \frac{B^2}{4} = O(\lambda_n^{\frac{1}{2}}). \end{aligned} \tag{120}$$

Combining (116),(117), Lemma 4, and (118) - (120), we find that for all sufficiently large n and $t \geq 1$,

$$\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2} \exp\left(-xt + cx^{\frac{1}{2}}\right) b_n(x)^{-1}dx$$

$$
\begin{aligned}
&\leq \quad \exp\big(c^2(2t)^{-1}\big)\Big((\frac{\pi}{\lambda_n t})^{\frac{1}{2}}\exp\big(-\frac{B^2}{4}t+\frac{1}{2}Bc+n^{-\frac{1}{3}}\big)+4\exp\big(-\frac{1}{2}\lambda_n t-\lambda_n^{\frac{1}{2}}t-\frac{B^2}{4}t\big)\Big)\\
&\leq \quad \exp\big(c^2(2t)^{-1}-\frac{B^2}{4}t+\frac{1}{2}Bc+n^{-\frac{1}{3}}\big)\Big((\frac{\pi}{\lambda_n t})^{\frac{1}{2}}+4\exp\big(-\frac{1}{2}\lambda_n t-\lambda_n^{\frac{1}{2}}t\big)\Big)\\
&\qquad \text{since } \exp\big(\frac{1}{2}Bc+n^{-\frac{1}{3}}\big)>1\\
&\leq \quad \exp\big(c^2(2t)^{-1}-\frac{B^2}{4}t+\frac{1}{2}Bc+n^{-\frac{1}{3}}\big)\Big((\frac{\pi}{\lambda_n t})^{\frac{1}{2}}+4(\frac{1}{2}\lambda_n t)^{-1}(\lambda_n^{\frac{1}{2}}t)^{-1}\Big)\\
&= \quad (\frac{\pi}{\lambda_n t})^{\frac{1}{2}}\exp\big(c^2(2t)^{-1}-\frac{B^2}{4}t+\frac{1}{2}Bc+n^{-\frac{1}{3}}\big)\Big(1+8\big(\pi^{\frac{1}{2}}\lambda_n t^{\frac{3}{2}}\big)^{-1}\Big)
\end{aligned}
$$

$$
\leq \quad (\frac{\pi}{\lambda_n t})^{\frac{1}{2}}\exp\big(c^2(2t)^{-1}-\frac{B^2}{4}t+\frac{1}{2}Bc+n^{-\frac{1}{3}}\big)\big(1+8\lambda_n^{-1}\big) \quad \text{since } t\geq 1, \pi^{\frac{1}{2}}\geq 1. \tag{121}
$$

Furthermore, since $1+8\lambda_n^{-1}=1+O(n^{-1})$, and $\exp(n^{-\frac{1}{3}})=1+O(n^{-\frac{1}{3}})$ by a simple Taylor series expansion, it follows that $(1+8\lambda_n^{-1})\exp(n^{-\frac{1}{3}})=1+O(n^{-\frac{1}{3}})$. Thus for all sufficiently large n, $(1+8\lambda_n^{-1})\exp(n^{-\frac{1}{3}})\leq 1+n^{-\frac{1}{4}}=q_n$. Combining with (121), and noting that in all cases 'for all sufficiently large n' can be defined in terms of C only (as opposed to c), completes the proof of the lemma. ☐

Proof of Theorem 7. Suppose $B\in(0,B^*)$, $a_1,a_2\in\mathbb{R}$. Let $\alpha=\max(B,|a_1|,|a_2|)$. Then for all sufficiently large n and all $t\geq 1$,

$$
\sqrt{n}|P^n_{\lceil n+a_1\sqrt{n}\rceil,\lceil n+a_2\sqrt{n}\rceil}(t)-P^n_{\lceil n+a_2\sqrt{n}\rceil}(\infty)| \tag{122}
$$

$$
\begin{aligned}
&\leq \quad (2\pi)^{-1}q_n\exp\big(B^2\big)(\lambda_n n^{\frac{1}{2}})^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt\big)|Q_{n,i}(x)|c_n(x)^{-\frac{1}{2}}|Q_{n,j}(x)|c_n(x)^{-\frac{1}{2}}b_n(x)dx\\
&\qquad \text{by Theorem 9, Corollary 1, and Lemma 5}\\
&\leq \quad (2\pi)^{-1}q_n\exp\big(B^2\big)(\lambda_n n^{\frac{1}{2}})^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt\big)h_n(x)^2\exp\big(2(\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)b_n(x)dx\\
&\qquad \text{by applying Lemma 32 to both } |Q_{n,i}(x)|c_n(x)^{-\frac{1}{2}} \text{ and } |Q_{n,j}(x)|c_n(x)^{-\frac{1}{2}}
\end{aligned}
$$

$$
= \quad 2\pi^{-1}q_n\exp\big(B^2\big)\frac{n^{\frac{3}{2}}}{\lambda_n}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt+2(\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)b_n(x)^{-1}dx. \tag{123}
$$

We now bound the integral appearing in (123), which is equivalent to

$$
\exp\big(2\alpha^2+2\alpha n^{-\frac{1}{5}}\big)\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2}\exp\big(-xt+6(\alpha+n^{-\frac{1}{5}})x^{\frac{1}{2}}\big)b_n(x)^{-1}dx
$$

$$
\leq q_n(\frac{\pi}{t\lambda_n})^{\frac{1}{2}}\exp\Big(2\alpha^2+2\alpha n^{-\frac{1}{5}}+3B(\alpha+n^{-\frac{1}{5}})+18(\alpha+n^{-\frac{1}{5}})^2t^{-1}-\frac{B^2}{4}t\Big) \tag{124}
$$

for all sufficiently large n, by applying Lemma 34 with $C=6(\alpha+1), c=6(\alpha+n^{-\frac{1}{5}})$.

Also, for all sufficiently large n,

$$\exp\big((2\alpha+3B)n^{-\frac{1}{5}}\big) = 1 + O(n^{-\frac{1}{5}}) \le 1 + n^{-\frac{1}{4}} = q_n \text{ by a simple Taylor series expansion,} \tag{125}$$

and similarly

$$\exp\big((36\alpha n^{-\frac{1}{5}} + 18n^{-\frac{2}{5}})t^{-1}\big) = 1 + O(n^{-\frac{1}{5}}) \le 1 + n^{-\frac{1}{4}} = q_n \text{ since } \ t \ge 1. \tag{126}$$

Combining (123), (125), and (126) we find that for all sufficiently large n, and all $t \ge 1$, the integral appearing in (123) is at most

$$q_n^3(\frac{\pi}{t\lambda_n})^{\frac{1}{2}} \exp\Big(2\alpha^2 + 3B\alpha + 18\alpha^2 t^{-1} - \frac{B^2}{4}t\Big). \tag{127}$$

Combining (123) and (127) we find that (122) is at most

$$2\pi^{-1} q_n \exp\big(B^2\big)\frac{n^{\frac{3}{2}}}{\lambda_n}\Bigg(q_n^3(\frac{\pi}{t\lambda_n})^{\frac{1}{2}} \exp\Big(2\alpha^2 + 3B\alpha + 18\alpha^2 t^{-1} - \frac{B^2}{4}t\Big)\Bigg)$$

$$\begin{aligned} &= 2(\pi t)^{-\frac{1}{2}} q_n^4 (\frac{n}{\lambda_n})^{\frac{3}{2}} \exp\Big(B^2 + 2\alpha^2 + 3B\alpha + 18\alpha^2 t^{-1} - \frac{B^2}{4}t\Big) \\ &\le 2(\pi t)^{-\frac{1}{2}} q_n^4 (\frac{n}{\lambda_n})^{\frac{3}{2}} \exp\Big(2(\alpha+B)^2 + 18\alpha^2 t^{-1} - \frac{B^2}{4}t\Big). \end{aligned} \tag{128}$$

Furthermore,

$$\frac{n}{\lambda_n} = 1 + \frac{Bn^{\frac{1}{2}}}{n - Bn^{\frac{1}{2}}} = 1 + O(n^{-\frac{1}{2}}) \le q_n \text{ for all sufficiently large n.} \tag{129}$$

The first part of Theorem 7 follows from (128) and (129), by noting that

$$q_n^{5.5} = 1 + O(n^{-\frac{1}{4}}) \le 1 + n^{-\frac{1}{8}} \text{ for all sufficiently large n.} \tag{130}$$

Similarly, for all sufficiently large n and all $t \ge 1$,

$$|P^n_{i,\le j}(t) - P^n_{\le j}(\infty)| \tag{131}$$

$$\leq \quad (2\pi)^{-1}q_n \exp\big(B^2\big)n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2} \exp\big(-xt\big)x^{-1}|Q_{n,i}(x)|c_n(x)^{-\frac{1}{2}}|Q_{n,j+1}(x)-Q_{n,j}(x)|c_n(x)^{-\frac{1}{2}}b_n(x)dx$$

by Lemma 33, Corollary 1, and Lemma 5

$$\leq \quad (2\pi)^{-1}q_n \exp\big(B^2\big)n^{-1}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2} h_n(x)^2 r_n(x)^{\frac{1}{2}}\exp\big(-xt+2(\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)x^{-1}b_n(x)dx$$

by applying Lemma 32 to both $|Q_{n,i}(x)|c_n(x)^{-\frac{1}{2}}$ and $|Q_{n,j+1}(x)-Q_{n,j}(x)|c_n(x)^{-\frac{1}{2}}$

$$= \quad 2\pi^{-1}q_n^{\frac{3}{2}}\exp\big(B^2\big)n^{\frac{1}{2}}\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2} \exp\big(-xt+2(\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)x^{-\frac{1}{2}}b_n(x)^{-1}dx$$

$$\leq \quad 2\pi^{-1}q_n^{\frac{3}{2}}\exp\big(B^2\big)n^{\frac{1}{2}}(2B^{-1})\int_{(\sqrt{n}-\sqrt{\lambda_n})^2}^{(\sqrt{n}+\sqrt{\lambda_n})^2} \exp\big(-xt+2(\alpha+n^{-\frac{1}{5}})(\alpha+3x^{\frac{1}{2}})\big)b_n(x)^{-1}dx$$

since $x^{-\frac{1}{2}} \leq \Big((\sqrt{n}-\sqrt{\lambda_n})^2\Big)^{-\frac{1}{2}} = (\sqrt{n}-\sqrt{\lambda_n})^{-1} \leq 2B^{-1}$ by Lemma 4

$$\leq \quad 4(B\pi)^{-1}q_n^{\frac{3}{2}}\exp\big(B^2\big)n^{\frac{1}{2}}\Bigg(q_n^3\big(\frac{\pi}{t\lambda_n}\big)^{\frac{1}{2}}\exp\Big(2\alpha^2+3B\alpha+18\alpha^2t^{-1}-\frac{B^2}{4}t\Big)\Bigg) \quad \text{By (127)}$$

$$= \quad 4q_n^{4.5}(B^2\pi t)^{-\frac{1}{2}}\big(\frac{n}{\lambda_n}\big)^{\frac{1}{2}}\exp\Big(B^2+2\alpha^2+3B\alpha+18\alpha^2t^{-1}-\frac{B^2}{4}t\Big).$$

The second part of Theorem 7 follows by combining the above with (129) and (130). □

8.1. *Comparison to other bounds from the literature.* In this subsection we compare the bounds from Theorem 7 to two other explicit bounds given in the literature [46],[7]. In both cases we will prove that the bounds from the literature (applied to $|P^n_{n,\leq n}(t)-P^n_{\leq n}(\infty)|$ for $0<B<B^*$) grow with n, rendering them impractical in the H-W regime. We begin with the bounds given in [46], which translate to the statement that for each $B \in (0,B^*)$ there exists N_B s.t. $n \geq N_B$ implies that for all $t \geq 0$,

(132)

$$|P^n_{n,\leq n}(t)-P^n_{\leq n}(\infty)| \leq 4(n-1)\Big(\sum_{i=1}^{\infty}\big((\frac{n}{n-1})^i-1\big)P^n_i(\infty)+\big((\frac{n}{n-1})^n-1\big)\big(1-2P^n_n(\infty)\big)\Big)e^{-\frac{B\sqrt{n-1}}{n-1}t}.$$

Note that since $\lim_{n\to\infty}\frac{B\sqrt{n-1}}{n-1}=0$, the exponential rate of convergence demonstrated by (132) goes to zero as $n\to\infty$, rendering the bound in [46] ineffective. We now examine the bounds given in [7], which translate to the statement that for each $B\in(0,B^*)$ there exists N_B s.t. $n\geq N_B$ implies that for all $t\geq 0$,

$$|P^n_{n,\leq n}(t)-P^n_{\leq n}(\infty)| \leq \big(P^n_n(\infty)^{-1}-1\big)^{\frac{1}{2}}e^{-\gamma_n t}. \tag{133}$$

It is well-known (see [22]) that $\liminf_{n\to\infty}\frac{(P^n_n(\infty)^{-1}-1)^{\frac{1}{2}}}{n^{\frac{1}{4}}}>0$. It follows that the prefactor demonstrated by (133) diverges as $n\to\infty$, rendering the bound in [7] ineffective.

It should be noted that although the bounds given in [46] and [7] are ineffective for the particular events of interest in this paper, both bounds hold in much greater generality, and thus remain interesting and applicable in a variety of other settings.

9. Conclusion and Open Questions. In this paper we studied the rate of convergence to stationarity of the $M/M/n$ queue in the H-W Regime. We explicitly computed the limit of the exponential rate of convergence to stationarity (*spectral gap*). We proved that there is an interesting *phase transition* in the system's behavior, occuring when the excess parameter B reaches $B^* \approx 1.85772$. For $B < B^*$, the exponential rate of convergence is $\frac{B^2}{4}$; above B^* it is the solution to an equation involving the parabolic cylinder functions. We showed that this transition asymptotically characterizes a phenomenon previously observed to occur for fixed n, unifying and simplifying several earlier lines of work. We also explicitly bounded the prefactor governing the rate of convergence uniformly over n (for $B < B^*$), and used our bounds to offer managerial insight into the operation of a call center.

This work leaves several interesting directions for future research. There are many open questions related to the interaction between weak convergence and convergence to stationarity. Although our results and those of [30] show that for the $M/M/n$ queue in the H-W regime there is an 'interchange of limits' in this regard, it is unknown to what extent such an interchange must hold in general. It would also be interesting to prove that a *phase transition* occurs in other models, such as queues with abandonments and non-Markovian service distributions.

10. Acknowledgements. The authors would like to thank Hans Blanc, Ton Dieker, Erik van Doorn, Peter Glynn, Johan van Leeuwaarden, Bill Massey, Josh Reed, and Ward Whitt for their helpful discussions and insights. The authors especially thank Johan van Leeuwaarden for his insights into the parabolic cylinder functions, and sharing an early draft of his work.

REFERENCES

[1] Z. Aksin, M. Armory, and V. Mehrotra, *The modern call center: a multi-disciplinary perspective on operations management research.*, Production and Operations Management **16** (2007), no. 6, 665–688.

[2] J.D. Atkinson and T.K. Caughley, *Spectral density of piecewise linear first order systems excited by white noise.*, Int. J. Non-Linear Mechanics (1968), no. 3, 137–156.

[3] H. Buchholz, *The confluent hypergeometric function.*, Springer-Verlag, New York, 1969.

[4] H. Callaert, *Exponential ergodicity for birth-death processes.*, Ph.D. Thesis, University of Louvain (1971).

[5] ______, *On the rate of convergence in birth-and-death processes.*, Bull.Soc.Math.Belg. **26** (1974), 173–184.

[6] M. Chen, *Exponential l^2 convergence and l^2 spectral gap for markov processes.*, Acta Mathematica Sinica, New Series **7** (1991), no. 1, 19–37.

[7] ______, *Estimate of exponential convergence rate in total variation by spectral gap.*, Acta Mathematica Sinica **14** (1998), no. 1, 9–16.

[8] T.S. Chihara, *An introduction to orthogonal polynomials.*, Gordon and Breach, New York, 1978.

[9] D. Chruscinski, *Quantum mechanics of damped systems. II. damping and parabolic potential barrier.*, Journal of Mathematical Physics **45** (2004), no. 3, 841–854.

[10] A. Clarke, *A waiting line process of markov type.*, The Annals of Mathematical Statistics **27** (1956), no. 2, 452–459.

[11] P. Diaconis, *The cutoff phenomenon in finite markov chains.*, Proc. Nat. Acad. Sci. (1996), no. 4, 1659–1664.

[12] D. Dominici, *Asymptotic analysis of the askey-scheme I: from Krawtchouk to Charlier.*, Cent. Eur. J. Math. **5** (2007), no. 2, 280–304.

[13] ______, *Asymptotic analysis of the Krawtchouk polynomials by the WKB method .*, Ramananujan J. **15** (2008), 303–338.

[14] T.M. Dunster, *Uniform asymptotic expansions for Charlier polynomials.*, J. Approx. Theory **112** (2001), no. 1, 93–133.

[15] A. Erdelyi, W. Magnus, F. Oberhettinger, F. Tricomi, and H. Bateman, *Higher transcendental functions, volume II*, McGraw-Hill Book Company, New York, 1953.

[16] A.K. Erlang, *On the rational determination of the number of circuits.*, The Copenhagen Telephone Company, Copenhagen, 1948.

[17] S. Ethier and T. Kurtz, *Markov processes: characterization and convergence*, Wiley and Sons, 2005.

[18] C. Fricker, P. Robert, and D. Tibi, *On the rates of convergence of Erlang's model.*, J. Appl. Probab. **36** (1999), 1167–1184.

[19] D. Gamarnik and P. Momcilovic, *Steady-state analysis of a multi-server queue in the Halfin-Whitt regime*, Advances in Applied Probability **40** (2008), 548–577.

[20] N. Gans, G. Koole, and A. Mandelbaum, *Telephone call centers: tutorial, review, and research prospects.*, Manufacturing and Service Operations Management **5** (2003), no. 2, 79–141.

[21] I.S. Gradshteyn and I.M.Ryzhik, *Table of integrals, series, and products.*, Academic Press, New York, 1980.

[22] S. Halfin and W. Whitt, *Heavy-traffic limits for queues with many exponential servers.*, Operations Research **29** (1981), no. 3, 567–588.

[23] E. Hille, *Analytic function theory*, Ginn and Company, 1959.

[24] D. Iglehart, *Limiting diffusion approximations for the many server queue and the repairman problem.*, Journal of Applied Probability **2** (1965), 429–441.

[25] D. Jagerman, *Some properties of the Erlang loss function.*, Bell System Techn. J. **53** (1974), no. 3, 525–551.

[26] P. Jelenkovic, A. Mandelbaum, and P. Momcilovic, *Heavy traffic limits for queues with many deterministic servers*, Queueing Systems: Theory and Applications **47** (2004), no. 1-2, 53–69.

[27] S. Karlin and J. McGregor, *The differential equation of birth-and-death processes, and the Stieltjes moment problem*, Trans. Amer. math. Soc. **85** (1957), 489–546.

[28] ———, *Many-server queueing processes with Poisson input and exponential service times.*, Pacific J. Math. **8** (1958), 87–118.

[29] M. Kijima, *Evaluation of the decay parameter for some specialized birth-death processes.*, J.Appl.Prob. **29** (1992), 781–791.

[30] C. Knessl and J. Leeuwaarden, *Transient behavior of the Halfin-Whitt diffusion*, Preprint (2008).

[31] W. Ledermann and G. Reuter, *Spectral theory for the differential equations of simple birth and death processes.*, Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences, **246** (1954), no. 914, 321–369.

[32] A. Mandelbaum and P. Momcilovic, *Queues with many servers: The virtual waiting-time process in the QED regime*, Math. Oper. Res. **33** (2008), no. 3, 561–586.

[33] P. Moorse, *Stochastic properties of waiting lines.*, Journal of the Operations Research Society of America **3** (1955), no. 3, 255–261.

[34] A. Puhalski and M. Reiman, *The multiclass GI/PH/N queue in the Halfin-Whitt regime.*, Advances in Applied Probability **32** (2000), no. 3, 564–595.

[35] R.Bo and R. Wong, *Uniform asymptotic expansion of Charlier polynomials.*, Methods Appl. Anal. **1** (1994), no. 3, 294–313.

[36] J. Reed, *The G/GI/n queue in the Halfin-Whitt regime.*, Annals of Applied Probability **19** (2009), no. 6, 2211–2269.

[37] T. Saaty, *Time-dependent solution of the many-server Poisson queue.*, Operations Research **8** (1960), no. 6, 755–772.

[38] E.A. van Doorn, *Stochastic monotonicity and queueing applications of birth death processes.*, Lecture Notes in Statist. **4** (1981).

[39] ———, *On oscillation properties and the interval of orthogonality of orthogonal polynomials.*, SIAM J. Math. Anal. **15** (1984), 1031–1042.

[40] ———, *Conditions for exponential ergodicity and bounds for the decay parameter of a birth-death process.*, Advances in Applied Probability **17** (1985), 514–530.

[41] ———, *Representations and bounds for zeros of orthogonal polynomials and eigenvalues of sign-symmetric tri-diagonal matrices.*, J.Approx.Theory **51** (1987), 254–266.

[42] E.A. van Doorn and A.I. Zeifman, *On the speed of convergence to stationarity of the Erlang loss system*, Queueing Systems **63** (2009), 241–252.

[43] E.A. van Doorn, A.I. Zeifman, and T.L. Panfilova, *Bounds and asymptotics for the rate of convergence of birth-death processes.*, Teor. Veroyatnost. I Primenen. **54** (2009), 18–38.

[44] Shisheng Xie and Charles Knessl, *On the transient behaviour of the Erlang loss model: heavy usage asymptotics*, SIAM J. Appl. Math. **53** (1993), no. 2, 555–599.

[45] R. Young, *Euler's constant*, The Mathematical Gazette (1991), no. 75, 187–190.

[46] A.I. Zeifman, *Some estimates of the rate of convergence for birth and death processes.*, Journal of Applied Probability **28** (1991), no. 2, 268–277.

Operations Research Center and Sloan School of Management, MIT, Cambridge, MA, 02139
E-mail: gamarnik@mit.edu; dag3141@mit.edu

This figure "zetaB_jpg.jpg" is available in "jpg" format from:

http://arxiv.org/ps/1003.2004v1